


Modern global carbon cycle and terrestrial/oceanic subsidies to anthropogenic emissions

C. David Keeling, 1928-2005




Dave Keeling (Scripps Institution of Oceanography, La Jolla, California) provided the longest continuous record of atmospheric CO₂ measurements.

In recognition of this work, Dr. Keeling was awarded the Medal of Science, the highest honor the U.S. awards a scientist.

Atmospheric CO₂ at Mauna Loa Observatory

Scripps Institution of Oceanography
NOAA Earth System Research Laboratory




http://www.esrl.noaa.gov/gmd/ccag/trends/mlo_full

$\text{CH}_2\text{O} + \text{O}_2 \longrightarrow \text{energy} + \text{H}_2\text{O} + \text{CO}_2$

Fossil fuels are the dominant source of energy in the United States, and in the industrialized world

Figure 1. U.S. primary energy consumption, 1980-2035

quadrillion Btu




<http://www.eia.doe.gov/oia/aeo/execssummary.html>

No significant changes in energy sources are expected in the next 25 years.


Three uses of energy dominate carbon dioxide emissions

Figure 4. U.S. energy-related carbon dioxide emissions, 2008 and 2035



<http://www.eia.doe.gov/oia/aeo/execssummary.html>

CO₂ emissions from energy use are estimated to increase by approximately 9 % between 2008 and 2035, which is an annual increase of 0.3 %.



Magnitude of carbon pools

Reservoir	Carbon Stored [Pg]
Sediments and rocks	60,000,000
Intermediate/deep ocean	38,000
Soils	1,500-2,350
Surface ocean	1,000
Atmosphere	760
Vegetation	650
Fossil fuels	3400


Terrestrial ecosystems are large carbon pools:
can potentially store a lot more.


Magnitude of carbon pool turnover times


Mean turnover time = pool size / input flux


Reservoir	Carbon Stored [Pg]
Soils	25-40
Surface ocean	11
Atmosphere	3-4
Vegetation	11
Fossil fuels	No input flux


Fossil fuels are not being created.















FACE experiments suggest plant growth can become nutrient limited at high CO₂ levels



Northern hemisphere terrestrial ecosystems are a major component of the current terrestrial ecosystem carbon sink

$$NEP = GPP - (R_p + R_a + R_m)$$



(Dragoni et al., 2011, *Global Change Biology*)


This data comes from direct measurements of ecosystem/atmosphere CO₂ exchange from eddy covariance, but there are multiple other lines of evidence that Northern hemisphere terrestrial ecosystems are major carbon sink (Tans and White, 1998, *Science*).

- ¹³C / ¹²C ratio in atmosphere.
- O₂ / N₂ ratio in atmosphere.
- Increase in CO₂ seasonal cycle amplitude.
- Climate variability and CO₂ increase rate.
- Forest inventories.
- Ocean inventories (especially ¹⁴C).

How much longer will terrestrial ecosystems in the Northern hemisphere act as carbon sinks?

Northern hemisphere terrestrial ecosystems are a major component of the current terrestrial ecosystem carbon sink

$$NEP = GPP - (R_p + R_a + R_m)$$




(Dragoni et al., 2011, *Global Change Biology*)

Possible causes of the Northern hemisphere carbon sink:


- Land use change (forest regrowth, woody encroachment, fire suppression).
- CO₂ fertilization (with no resource limitations, rising CO₂ stimulates photosynthesis).
- N fertilization (N deposition from pollution can enhance photosynthesis).
- Temperature (warming influences growing seasons and decomposition in soil).
- Water (if water is limiting, increased precipitation enhances photosynthesis).

Why are terrestrial ecosystems in the Northern hemisphere acting as carbon sinks?


Kyoto Protocol: Reduce greenhouse gas emissions to 5 % below the 1990 emission level



Projected changes in the global carbon cycle are highly dependent on future population growth and energy sources; high uncertainty



Consider the vulnerability of carbon pools in the 21st Century to temperature change: stored peats, increased fires, CH₄ hydrates



Global carbon cycle has been altered by “modern” CO₂ emissions

- Humans emit 2-3 times more CO₂ (from combustion of fossil fuel energy) in a year than the global terrestrial biosphere can store in a year.
- Is this energy consumption sustainable?
- Do we really need to use all this energy?
- Can the current level of fossil fuel energy usage be replaced with solar, wind, hydroelectric, and biofuel sources?
