

## OXYGEN ISOTOPE RATIOS OF WATERS AND RESPIRED CO<sub>2</sub> IN AMAZONIAN FOREST AND PASTURE ECOSYSTEMS

J. P. H. OMETTO,<sup>1,2</sup> L. B. FLANAGAN,<sup>3</sup> L. A. MARTINELLI,<sup>1</sup> AND J. R. EHLERINGER<sup>2,4</sup>

<sup>1</sup>Centro de Energia Nuclear na Agricultura, Avenida Centenário 303, Cep 13416-000, Piracicaba SP, Brazil

<sup>2</sup>Stable Isotope Ratio Facility for Environmental Research, Department of Biology, University of Utah,  
257 South 1400 East, Salt Lake City, Utah 84112-0840 USA

<sup>3</sup>Department of Biological Sciences, University of Lethbridge, 4401 University Drive, Lethbridge,  
Alberta T1K 3M4, Canada

**Abstract.** The oxygen isotope ratio ( $\delta^{18}\text{O}$ , SMOW) of atmospheric CO<sub>2</sub> is a powerful indicator of large-scale CO<sub>2</sub> exchange on land. Oxygen isotopic exchange between CO<sub>2</sub> and water in leaves and soils controls the  $\delta^{18}\text{O}$  of atmospheric CO<sub>2</sub>. Currently there is little empirical information on the spatial and temporal variation in the  $\delta^{18}\text{O}$  of leaf and stem water in tropical ecosystems. We measured the seasonal dynamics of  $\delta^{18}\text{O}$  in atmospheric CO<sub>2</sub> and water in different ecosystem compartments in both primary forest and pasture ecosystems in three different regions of the Amazonian Basin of Brazil (Ji-Paraná, Manaus, and Santarém). Within regions, the source (stem) water  $\delta^{18}\text{O}$  values for primary forests and pastures were similar; neither vegetation type exhibited distinct wet–dry season patterns. Daytime leaf water isotope ratios were strongly correlated with predictions of the Craig-Gordon model. The  $\delta^{18}\text{O}$  value of leaf water was positively correlated with leaf height above ground because of associated variation in vapor pressure deficit and the  $\delta^{18}\text{O}$  of atmospheric water vapor within forest canopies. Consistent with these observations, the  $\delta^{18}\text{O}$  value of leaf cellulose was positively correlated with forest height. Leaf water from pasture grasses was more <sup>18</sup>O enriched than leaf water from forest vegetation. There was a tendency for daytime leaf water to be more enriched in <sup>18</sup>O during the dry season, reflecting generally lower humidity conditions during the dry season. Nighttime measurements of the oxygen isotope ratio of ecosystem respired CO<sub>2</sub> in both forest and pasture vegetation were not consistent with values expected for CO<sub>2</sub> in equilibrium with stem (soil) water, despite nighttime vapor pressure deficits close to zero. Apparently, the  $\delta^{18}\text{O}$  of leaf water lagged and did not attain isotopic equilibrium at night. Thus, the deviation of nighttime  $\delta^{18}\text{O}$  values of ecosystem respiration from that expected from a CO<sub>2</sub> efflux in equilibrium with soil (stem) water increased as  $\delta^{18}\text{O}$  values of ecosystem respiration became <sup>18</sup>O enriched. Discrimination against CO<sub>2</sub> containing <sup>18</sup>O ( $\Delta\text{C}^{18}\text{OO}$ ) during photosynthesis was calculated based on measured leaf water  $\delta^{18}\text{O}$  values. Forests had consistently higher modeled  $\Delta\text{C}^{18}\text{OO}$  values than pastures. The daytime isotope effects we calculate for photosynthesis and respiration were consistent with previous model predictions of a strong depletion of <sup>18</sup>O in atmospheric CO<sub>2</sub> over the Amazon Basin of Brazil.

**Key words:** carbon cycle; carbon dioxide (CO<sub>2</sub>); oxygen isotope ratio; photosynthesis; respiration; stable isotope ratio; tropical forests.

### INTRODUCTION

The annual increase in atmospheric carbon dioxide concentration varies substantially from year to year, despite relatively constant anthropogenic CO<sub>2</sub> emissions from fossil fuel burning (Marland et al. 1999). This has led to the hypothesis that environmentally induced variation in carbon dioxide exchange in terrestrial ecosystems is largely responsible for the observed interannual variation in the rise of atmospheric CO<sub>2</sub> (Fung 2000). Measurement and analysis of the stable carbon isotope composition of atmospheric CO<sub>2</sub>

also support large yearly variation in terrestrial net ecosystem carbon uptake (Tans and White 1998). The mechanisms responsible for annual changes in terrestrial ecosystem CO<sub>2</sub> exchange are not well understood, but El Niño/La Niña events contribute substantially to yearly variation in temperature and precipitation with likely significant effects on ecosystem function in tropical regions (Tian et al. 1998, Behrenfeld et al. 2001). Strong interest has been expressed in using additional analyses of the concentration and stable isotope ratio of atmospheric carbon dioxide to infer changes in large-scale biosphere activity and to increase understanding of the controlling processes (Canadell et al. 2000).

The oxygen isotope ratio ( $\delta^{18}\text{O}$ ) of atmospheric CO<sub>2</sub> is a powerful indicator of large-scale CO<sub>2</sub> exchange on land (Farquhar et al. 1993, Ciais et al. 1997a, b, Gillon

Manuscript received 12 February 2003; revised 14 January 2004; accepted 15 January 2004; final version received 7 June 2004. Corresponding Editor: A. H. Goldstein.

<sup>4</sup> Corresponding author.

E-mail: ehleringer@biology.utah.edu

and Yakir 2001, Ehleringer et al. 2002). The primary factor influencing the  $^{18}\text{O}$  content of atmospheric  $\text{CO}_2$  is an equilibrium isotope effect that occurs between oxygen in water and oxygen in  $\text{CO}_2$  dissolved in the water. During photosynthesis,  $\text{CO}_2\text{-H}_2\text{O}$  exchange in the chloroplast tends to enrich the atmosphere in  $^{18}\text{O}$  because chloroplast water has a higher  $^{18}\text{O}:^{16}\text{O}$  ratio than water in the soil. Isotopic fractionation during transpiration causes the water pool remaining in leaves to become enriched in  $^{18}\text{O}$  relative to water taken up from the soil (Helliker and Ehleringer 2000, Yakir and Sternberg 2000). As a consequence, soil respiration releases  $\text{CO}_2$  to the atmosphere that is relatively  $^{18}\text{O}$  depleted, in contrast to the normal  $^{18}\text{O}$  enrichment associated with  $\text{CO}_2$  exchange during photosynthesis.

Seasonal changes in the  $\delta^{18}\text{O}$  value of atmospheric  $\text{CO}_2$  at both regional and global scales are largely determined by gross terrestrial fluxes of photosynthesis and respiration (Ciais and Meijer 1998). Francey and Tans (1987) and Friedli et al. (1987) first proposed that isotopic exchange involving soil and leaf water compartments in terrestrial ecosystems were the driving forces behind the observed latitudinal gradient in  $\delta^{18}\text{O}$  of  $\text{CO}_2$ . Subsequently, mechanistic models have been developed for the major isotopic fractionation processes contributing to variations in the  $\delta^{18}\text{O}$  value of atmospheric  $\text{CO}_2$  (Farquhar et al. 1993, Ciais et al. 1997a, b, Peylin et al. 1999, Riley et al. 2002). However, the full application of these models is difficult because of the complexity and interacting nature of the many controlling factors. In addition, there are limited field data available to validate major parameters in the models, and there is no information on seasonal variation and the effect of El Niño/La Niña cycles in tropical areas. Ciais and Meijer (1998) have specifically emphasized that the following information is necessary to reduce uncertainties in the application of  $\delta^{18}\text{O}$  in global carbon budget models. First, we need further assessment of the validity of the Craig-Gordon (1965) model (Flanagan et al. 1991) for calculating leaf water isotopic enrichment under field conditions. Second, better characterization is needed of the isotope ratio of ecosystem respired  $\text{CO}_2$  and its controlling factors. Despite the fact that the largest photosynthetic and respiratory fluxes are found in tropical ecosystems (Cramer et al. 1999, 2001), only a limited number of data sets exist that can be used to test model predictions in these ecosystems.

Our objective in this study was to measure seasonal and interannual variation in the  $\delta^{18}\text{O}$  value of leaf and stem (soil) water, and to compare these to measurements of the oxygen isotope ratio of ecosystem respired  $\text{CO}_2$ . We made measurements in primary forests and pastures in the Amazon Basin of Brazil in order to evaluate temporal changes in environmental conditions and shifts in land use for their effects on the oxygen isotope ratio of atmospheric  $\text{CO}_2$ . This research was part of a larger study, the Large Scale Biosphere At-

mosphere Experiment in Amazonia (LBA), which aims to improve our understanding of the interactions between the atmosphere and ecosystems of the Amazon Basin, a globally important biome.

#### Oxygen isotope ratio of ecosystem $\text{H}_2\text{O}$ and $\text{CO}_2$

Throughout this paper we use the conventional “delta” notation, which expresses the isotopic composition of a material relative to that of a standard:

$$\delta = \left( \frac{R_{\text{Sample}}}{R_{\text{Standard}}} - 1 \right) \quad (1)$$

where  $\delta$  is the isotope ratio and  $R$  is the molar ratio of heavy to light isotopes. The  $\delta$  values are conveniently expressed in parts per thousand (‰). The international standard for oxygen in water, cellulose, and gaseous forms is Standard Mean Ocean Water (SMOW).  $R_{\text{SMOW}}(^{18}\text{O}:^{16}\text{O}) = 2.0052 \times 10^{-3}$ .

In order to determine the impact of ecosystem gas exchange on the  $\delta^{18}\text{O}$  of atmospheric  $\text{CO}_2$ , we measured and/or calculated (1) the  $\delta^{18}\text{O}$  of leaf water over a diurnal cycle, (2) photosynthetic discrimination against  $\text{CO}_2$  molecules containing  $^{18}\text{O}$  ( $\Delta\text{C}^{18}\text{OO}$ ), and (3) the isotope ratio of  $\text{CO}_2$  respired at night from the entire ecosystem ( $\delta^{18}\text{O}_R$ ) and from the soil ( $\delta^{18}\text{O}_{R\text{-soil}}$ ). All  $\delta^{18}\text{O}_R$  values were calculated as the intercept from a Keeling plot, two-end-member mixing model (Pataki et al. 2003). Data for these analyses were from flask collections during brief nighttime periods (2–3 h following sunset) and are only presented when the statistical significance of the linear regression was  $P < 0.05$ .

The  $\delta^{18}\text{O}$  value of water in the roots and stems ( $\delta^{18}\text{O}_{\text{Stem}}$ ) remains the same as the source soil water until reaching the leaves, where evaporative enrichment occurs. Leaf water ( $R_{\text{Leaf}}$  or  $\delta^{18}\text{O}_{\text{Leaf}}$ ) can be calculated using a Craig-Gordon (1965) model as modified by Flanagan et al. (1991):

$$R_{\text{Leaf}} = \alpha^* \left[ \alpha_k R_{\text{Stem}} \left( \frac{e_i - e_a}{e_i} \right) + R_{\text{Atm}} \left( \frac{e_a}{e_i} \right) \right] \quad (2)$$

$$\delta^{18}\text{O}_{\text{Leaf}} = \left( \frac{R_{\text{Leaf}}}{R_{\text{SMOW}}} - 1 \right) \quad (3)$$

where  $R$  is the molar ratio of heavy to light isotopes (subscripts Atm, Leaf, and Stem refer to atmospheric water vapor, leaf water, and stem water, respectively) and  $e$  is the vapor pressure (subscripts a and i refer to air outside of a leaf and leaf intercellular air spaces, respectively). The liquid–vapor equilibrium fractionation factor is represented by  $\alpha^*$  ( $\alpha^* = 1.009$  at  $25^\circ\text{C}$ ) and varies with temperature according to the equations of Majoube (1971);  $\alpha_k$  is the kinetic fractionation associated with  $\text{H}_2\text{O}$  diffusion in air ( $\alpha_k = 1.0285$ ).

During photosynthetic gas exchange, discrimination against  $\text{CO}_2$  molecules containing  $^{18}\text{O}$  is influenced by diffusion and by oxygen isotope exchange with leaf

water. A portion of the CO<sub>2</sub> that enters a leaf during photosynthesis diffuses back to the atmosphere with an altered oxygen isotope ratio, after equilibration with leaf (chloroplast) water. The oxygen isotope exchange process is facilitated by the enzyme carbon anhydrase (CA), which rapidly catalyses the hydration of CO<sub>2</sub> and dehydration of HCO<sub>3</sub><sup>-</sup> in the chloroplast. In C<sub>3</sub> shrubs and trees CA activity is high, while CA activity is low in C<sub>4</sub> grasses (Gillon and Yakir 2001). Discrimination against C<sup>18</sup>O during photosynthetic gas exchange (ΔC<sup>18</sup>O) also depends on the gradient in concentration of CO<sub>2</sub> between the air and the chloroplast surface. These processes are all included in the model developed by Farquhar et al. (1993) and Farquhar and Lloyd (1993):

$$\Delta C^{18}O = \bar{a} + \left( \frac{c_c}{c_a - c_c} \right) \times \left[ \theta(\delta_c - \delta_a) - \left( \frac{\bar{a}}{\frac{c_c}{c_a - c_c} + 1} \right) (1 - \theta) \right] \quad (4)$$

$$\delta_c = \delta^{18}O_{\text{Leaf}} + \varepsilon \quad (5)$$

where  $c_c$  and  $c_a$  are the chloroplast and ambient air concentrations of CO<sub>2</sub>;  $\delta_c$  and  $\delta_a$  are  $\delta^{18}O$  values of CO<sub>2</sub> in the chloroplast and ambient air, respectively;  $\theta$  is the proportional extent of isotopic equilibrium between oxygen in CO<sub>2</sub> and oxygen in chloroplast water ( $\theta = 0.93$  for C<sub>3</sub> trees and shrubs and  $\theta = 0.3$  for C<sub>4</sub> grasses (Gillon and Yakir 2001));  $\varepsilon$  is the equilibrium fractionation between oxygen in CO<sub>2</sub> and oxygen in water (41.2‰ at 25°C; oxygen in CO<sub>2</sub> becomes enriched in <sup>18</sup>O relative to the oxygen in the water pool); and  $\bar{a}$  is the weighted mean fractionation during diffusion of CO<sub>2</sub> between the atmosphere and the chloroplast surface (Farquhar et al. 1993). We used measurements of leaf carbon isotope composition in the C<sub>3</sub> forest vegetation to estimate  $c_c$  based on  $c_a$  values of 370 μmol of CO<sub>2</sub> per mol of air. The average  $\delta^{13}C$  value of “sun” leaves was approximately -30‰ in these forests (Ometto et al. 2002), which translates into a  $c_c$  value of 282 μmol/mol (Farquhar et al. 1989). For C<sub>4</sub> plants we assumed a  $c_c$  value of 180 μmol/mol (Farquhar et al. 1989). The  $\delta_a$  was assumed to be 41‰ for the model calculations.

The  $\delta^{18}O$  value of CO<sub>2</sub> respired from soils is controlled by isotopic fractionation during equilibration of soil CO<sub>2</sub> with water in the soil, and fractionation that occurs during diffusion of CO<sub>2</sub> out of the soil. Therefore, we calculate the  $\delta^{18}O$  of soil CO<sub>2</sub> efflux ( $\delta^{18}O_{\text{R-soil}}$ ) as

$$\delta^{18}O_{\text{R-soil}} = \delta^{18}O_{\text{Stem}} + \varepsilon - a \quad (6)$$

where  $a$  is assumed to have a value of 8.8‰ (Tans 1998).

## MATERIALS AND METHODS

### Study sites and precipitation patterns

The sites were located within the Amazon Basin near Santarém, Manaus, and Ji-Paraná, Brazil. In each re-

gion, measurements were collected at forest and pasture sites. Each of the forest sites were primary evergreen forests (“terra firme”) with a mean canopy height of 30–35 m, although some emergent trees reached 45–50 m. The major land use change replacing the primary forests in Amazonia is pastures, dominated by *Brachiaria* spp., a non-native C<sub>4</sub> grass widespread throughout the region.

The primary forest site near Santarém was located 67 km south of the city, in an area between the Tapajós River and the highway BR 163 (2.85° S, 54.05° W). The forest covers an area of ~600 000 ha and was established as a national forest in 1974 (FLONA Tapajós). The Santarém pasture site was located 77 km south of Santarém (3.02° S, 54.06° W). The Manaus forest site was located 70 km from the city (2.69° S, 60.11° W), in a forest reserve controlled by the Instituto Nacional de Pesquisas Amazonicas (INPA). The Manaus pasture site was located about 60 km from the forest site, along a secondary road (ZF-3). The grass at this site had less vigorous growth than the Santarém pasture grass. The soils at the Manaus and Santarém sites were deeply weathered oxisols (Hapludox) with high clay content (60–80%), low pH (4.0–4.3), and low nutrient content. The Ji-Paraná primary forest site (Rebio Jaru, 10.08° S, 61.92° W) was a reserve controlled by the Brazilian Environment Protection Agency (IBAMA), located north of Ji-Paraná (~80 km), by the Machado River (120 m above sea level). The pasture site (Fazenda Nossa Senhora da Aparecida; 10.75° S, 62.37° W) was a cattle ranch 50 km northeast from Ji-Paraná. The soil in both Ji-Paraná sites has been classified by Hodnett et al. (1996) as an orthic Acrisol, with 85%, or more, of sand at the surface layer. All sites, except the Manaus pasture, were official LBA-Ecology study areas. Field measurements were made during several field campaigns from March 1999 through to March 2001 at the Santarém and Manaus sites. Less frequent sampling occurred in Ji-Paraná.

At each of our study sites, there was a clear dry season (<100 mm precipitation per month), although the length and timing of the dry season was different among sites (Fig. 1). The transition from wet season to dry season (and vice versa) was not abrupt and did show some interannual variation. The monthly wet season precipitation was ~250–400 mm (Fig. 1).

### Water collection and analysis

Water samples were collected to characterize the stable isotope ratios of different water compartments in pasture and forest ecosystems. Xylem water samples were collected from stems ( $\delta^{18}O_{\text{Stem}}$ ) to determine plant water sources in the soil. Leaves were sampled for their leaf water ( $\delta^{18}O_{\text{Leaf}}$ ) and atmospheric water vapor ( $\delta^{18}O_{\text{Atm}}$ ) was trapped so that  $\delta^{18}O_{\text{Leaf}}$  values could be modeled. The leaves and stems were collected from several heights within the forest canopy, placed immediately inside individual glass vials sealed with a




FIG. 1. Mean monthly precipitation for Manaus, Santarém, and Ji-Paraná. The means are for the years 1968–1998 at Santarém, 1961–1990 at Manaus, and 1977–1997 at Ji-Paraná.

rubber stopper and wrapped with Parafilm. The samples were kept cold (0–5°C) in the field and later frozen in the lab until water was extracted using cryogenic vacuum distillation (Ehleringer et al. 2001). Atmospheric water vapor was collected by separately pumping air from different heights within the canopy (three in the forest and two in the pasture) through cold traps (dry ice–ethanol ice slurry) at a flow rate of 1 mL/s (Helliker et al. 2002). Oxygen isotope ratios for all water samples were determined by equilibration of water with CO<sub>2</sub> (Socki et al. 1999). Measurements of the oxygen isotope composition of the equilibrated CO<sub>2</sub> were performed by online injection of CO<sub>2</sub> into a coupled gas chromatograph and isotope ratio mass spectrometer (Delta Plus, Finnigan MAT, Bremen, Germany), operating in a continuous flow mode (Fessenden et al. 2002).

#### Air sample collection and analysis

Air samples from within the forest canopy and pasture sites were collected using Dekoron tubing (6 mm outside diameter) attached to a canopy access tower at different heights in the forest (42–31, 21, 7, 0.2 m with a variable inlet at the highest level dependent on the site) and attached to a pole at pasture (0.7, 0.2 m; Ometto et al. 2002). Air was pulled through the tubing using a battery operated pump (Capex V2X, Charles Austin, West Byfleet, Surrey, UK) at a rate of ~20 mL/s. As air flowed through the tubing, it was first dried by passing through a desiccant tube containing magnesium perchlorate and then through a 100-mL glass flask with Teflon stopcocks at either end; the pump was located at the end of this line (Ehleringer and Cook 1998). Flasks were flushed for 7–10 min before stopcocks on the flask were closed.

We sampled the CO<sub>2</sub> respired from soils using a closed cuvette system (Flanagan et al. 1999). A 50 × 40 × 40 cm cuvette was sealed to the soil surface by insertion 5 cm into the soil. A small fan operating at a low flow rate was used to mix air within this cuvette.

A long narrow tube was connected between the cuvette and the outside atmosphere to limit pressure differences from developing between inside and outside the cuvette. Air was moved slowly from the cuvette by a separate closed loop through an infrared gas analyzer (LiCor Gas Hound, LI-800, Lincoln, Nebraska, USA) to monitor CO<sub>2</sub> concentration within the cuvette. Attached to the side of the cuvette was a coupled desiccant trap and evacuated 100-mL flask. The CO<sub>2</sub> level was allowed to rise within the cuvette in response to soil respiration and at prescribed CO<sub>2</sub> intervals cuvette air was sampled using the attached flasks.

Stable isotope ratios of CO<sub>2</sub> in air samples collected in 100-mL glass flasks were analyzed using a coupled gas preconcentrator (PreCon) and isotope ratio mass spectrometer (Model 252, Finnigan MAT, Bremen, Germany) operating in a continuous flow mode. A 400-μL subsample of air from the flask was injected into a PreCon system, which allowed isolation of the CO<sub>2</sub> from all other gases before entering into the mass spectrometer. The long-term δ<sup>18</sup>O measurement precision of this technique has been ±0.23‰. After stable isotope ratio analysis of the flask had been completed, the CO<sub>2</sub> concentration was determined using a coupled bellows–infrared gas analyzer system (Bowling et al. 2001). The long-term [CO<sub>2</sub>] measurement precision has been ±0.3 ppm.

We used a mixing model developed by Keeling (1958), and applied extensively in studies of ecosystem-scale gas exchange (Buchmann et al. 1997, Flanagan and Ehleringer 1998, Bowling et al. 2002, Fessenden and Ehleringer 2002, Ometto et al. 2002, Pataki et al. 2003), to calculate the isotope ratio of CO<sub>2</sub> respired by an entire ecosystem (δ<sup>18</sup>O<sub>R</sub>) and that released from the soil surface (δ<sup>18</sup>O<sub>R-soil</sub>). Estimates for δ<sup>18</sup>O<sub>R</sub> were obtained from the y-intercept of a linear regression between δ<sub>a</sub> and c<sub>a</sub><sup>-1</sup> values measured on air samples collected at different heights within a vegetation canopy at night when photosynthesis was not active. This analysis was also applied to the air samples collected from the soil respiration chamber. The δ<sup>18</sup>O<sub>R</sub> values presented were all statistically significant (*P* < 0.05). It is important to recognize that the isotope ratio of soil respired CO<sub>2</sub> (δ<sup>18</sup>O<sub>R-soil</sub>), as calculated by Eq. 6, is only one component of the isotope ratio of CO<sub>2</sub> respired by an entire ecosystem (represented by δ<sup>18</sup>O<sub>R</sub>).

The original data have been archived within Ecological Archives as a Supplement and additionally at NASA LBA-ECO and CPTEC LBA.<sup>5,6</sup>

## RESULTS

### Stem and leaf water δ<sup>18</sup>O values

Over a 24-mo period, stem water δ<sup>18</sup>O values varied from –9‰ to –3‰ across the three study locations (Fig. 2). Within a region, the δ<sup>18</sup>O<sub>Stem</sub> values of plants

<sup>5</sup> <http://beija-flor.ornl.gov/lba/>

<sup>6</sup> <http://lba.cptec.inpe.br>




FIG. 2. Temporal variation (mean  $\pm$  1 SE) in the oxygen isotope ratio ( $\delta^{18}\text{O}$ , ‰) of stem and leaf water in forest and pasture sites in Brazil.

at forest and pasture sites varied in concert and were very similar on any given sampling date. There was no clear indication of any seasonal (wet vs. dry season) pattern in  $\delta^{18}\text{O}_{\text{Stem}}$  values within a site at any of the six different locations. At the Manaus and Santarém sites where data collections were most extensive (including both pasture and forest), both the minimum and maximum  $\delta^{18}\text{O}_{\text{Stem}}$  values occurred during the wet season period.

While the  $\delta^{18}\text{O}$  value of stem water remained relatively constant, the  $\delta^{18}\text{O}_{\text{Leaf}}$  values were enriched in  $^{18}\text{O}$  relative to stem water and showed much larger temporal and spatial variation (Figs. 2, 3). We used air temperature and humidity data, along with measurements of

$\delta^{18}\text{O}_{\text{Stem}}$  and  $\delta^{18}\text{O}_{\text{Atm}}$  values to model the expected variation in the oxygen isotope ratio of leaf water. At the start of our study, continuous measurements of air temperature and humidity were only available at one study location, the Manaus primary forest site. We observed a strong agreement ( $r^2 = 0.83$ ,  $P < 0.01$ ) between modeled and observed leaf water  $\delta^{18}\text{O}$  values at the Manaus forest site (Fig. 3). Since  $\delta^{18}\text{O}_{\text{Stem}}$  values were relatively constant, the Craig-Gordon (1965) model (Eq. 2) predicts that leaf water should vary in response to changes in vapor pressure deficit (VPD) and  $\delta^{18}\text{O}_{\text{Atm}}$  (Flanagan et al. 1991). Variation in VPD, in particular, might be expected to result in differences in the  $\delta^{18}\text{O}$  value of leaf water (1) along a height gradient within a forest canopy, (2) between wet and dry seasons, and (3) in a comparison of forest and pasture sites. We used our measurements and model calculations to examine the influence of these three factors on the oxygen isotope ratio of leaf water.

There was a consistent vertical gradient in the mid-day  $\delta^{18}\text{O}_{\text{Leaf}}$  values in the forest canopies (Fig. 4). Upper canopy leaves from forest sites had water that was more enriched in  $^{18}\text{O}$  by 4–6‰ than leaves from lower positions in the canopy. This vertical gradient in leaf water isotope ratio was likely caused by changes in both  $\delta^{18}\text{O}_{\text{Atm}}$  values and VPD throughout the forest canopy. At the Santarém forest site, there was also a strong vertical gradient for the  $\delta^{18}\text{O}$  value of cellulose purified from leaves (Fig. 5). The oxygen isotope composition of leaf cellulose represents a temporal integration of the  $\delta^{18}\text{O}$  of leaf water present during cellulose synthesis (Roden et al. 2000).




FIG. 3. The relationship between measured and modeled oxygen isotope ratio ( $\delta^{18}\text{O}$ , ‰) of leaf water in the Manaus F2 forest in Brazil.




FIG. 4. The relationship between leaf height and the oxygen isotope ratio of leaf water (mean  $\pm$  1 SE) in forest sites in Brazil. Data from the wet season (upper panels: >100 mm precipitation/mo) and dry season (lower panels: <100 mm precipitation/mo) are shown in separate panels for each site.

Forest leaf water  $\delta^{18}\text{O}$  values tended to be more enriched in  $^{18}\text{O}$  during the dry season than the wet season (Fig. 4). Our sampling was too limited to rigorously test for dry vs. wet season differences in pasture ecosystems.

Leaf water from pasture grasses was generally more enriched in  $^{18}\text{O}$  during the day than leaf water from forest vegetation, with  $\delta^{18}\text{O}$  values ranging from 1.3‰ to 15.2‰ in Manaus pasture, and similar values occurring in pastures from other regions (Fig. 2). Model calculations, for a comparison of leaf water  $\delta^{18}\text{O}$  values in forest and pasture sites in Santarém during September 2000, showed that greater enrichment of  $^{18}\text{O}$  in leaf water was expected in the pasture site (Fig. 6). This




FIG. 5. The relationship between leaf height and the oxygen isotope ratio of leaf cellulose (mean  $\pm$  1 SE) in the Santarém primary forest site in Brazil.

difference in modeled  $\delta^{18}\text{O}_{\text{Leaf}}$  was most likely a result of the higher VPD in the pasture site, because measurements of  $\delta^{18}\text{O}_{\text{Atm}}$  and  $\delta^{18}\text{O}_{\text{Stem}}$  were very similar in the forest and pasture sites (Fig. 6).

#### *δ<sup>18</sup>O of respired CO<sub>2</sub> in forest and pasture ecosystems*

Highly significant linear relationships occurred between the inverse of  $\text{CO}_2$  concentration and the  $\delta^{18}\text{O}$  values of  $\text{CO}_2$  measured on air samples collected within a vegetation canopy at night ( $r^2 > 0.85$  typically; see Supplement) and in the soil respiration chamber (Fig. 7). In March 2001, the  $\delta^{18}\text{O}_{\text{Stem}}$  value for trees in the Santarém forest was  $-7\text{‰}$ . Therefore, we predicted (from Eq. 6) that  $\delta^{18}\text{O}_{\text{R-soil}}$  should be  $\sim 25\text{‰}$ , which is what was observed (Fig. 7). We repeated these  $\delta^{18}\text{O}_{\text{R-soil}}$  measurements on several occasions and each time the observed  $\delta^{18}\text{O}_{\text{R-soil}}$  values are within 1–2‰ of that predicted from measured  $\delta^{18}\text{O}_{\text{Stem}}$  values and Eq. 6.

At the whole ecosystem level, the oxygen isotope composition of nighttime respiration ( $\delta^{18}\text{O}_{\text{R}}$ ) should reflect the integrated inputs of  $\text{CO}_2$  respired by both aboveground vegetation and soil components. We observed temporal variation in  $\delta^{18}\text{O}_{\text{R}}$  values in both forest and pasture sites (Fig. 8). Forest ecosystem  $\delta^{18}\text{O}_{\text{R}}$  values ranged between 19.8‰ and 33.1‰, whereas  $\delta^{18}\text{O}_{\text{R}}$  values from pasture sites ranged between 22.8‰ and 39.2‰. In general,  $\delta^{18}\text{O}_{\text{R}}$  values were more  $^{18}\text{O}$  enriched during the dry season than during the wet season at both forest and pasture sites (Fig. 8). Some of the temporal variation in  $\delta^{18}\text{O}_{\text{R}}$  values was correlated with




FIG. 6. Comparison of modeled oxygen isotope ratio of leaf water in Santarém forest and pasture sites during September 2000. The left-hand panels represent the mean diurnal pattern for the 10-d period shown in the right-hand panels. The calculations used air temperature and humidity data collected by D. Fitzgerald at the LBA pasture site, and similar data collected by M. Goulden at the forest site. Measurements of the following values were also used in the calculations:  $\delta^{18}\text{O}_{\text{Stem}} = -3.50\text{‰}$  (forest) and  $-2.81\text{‰}$  (pasture);  $\delta^{18}\text{O}_{\text{Atm}} = -9.69\text{‰}$  (forest) and  $-8.51\text{‰}$  (pasture).

changes in the isotope ratio of source water as reflected in the  $\delta^{18}\text{O}_{\text{Stem}}$  values (Fig. 8).

We can interpret some of the variation in  $\delta^{18}\text{O}_{\text{R}}$  values, although we are restricted from a full interpretation because nighttime  $\delta^{18}\text{O}_{\text{Leaf}}$  values were not measured in this study. In these tropical ecosystems, nighttime humidity levels generally approached 100% shortly after sundown. For an initial interpretation of  $\delta^{18}\text{O}_{\text{R}}$  values, let us assume that the  $\delta^{18}\text{O}$  of leaf and soil water should

be similar at night. If so, then  $\delta^{18}\text{O}_{\text{Leaf}}$  values should approach that of  $\delta^{18}\text{O}_{\text{Stem}}$ , which we expect to be equivalent to  $\delta^{18}\text{O}$  values of the soil. Here we also assume that the  $\delta^{18}\text{O}$  values of soil water in the upper soil layers and stem water are identical. This is likely to hold in most cases, since these sites are extremely moist and the high precipitation inputs and well-shaded forest floor are likely not to have experienced much evaporative  $^{18}\text{O}$  enrichment. When interpreting  $\delta^{18}\text{O}_{\text{R}}$  values, Bowling et al. (2003) caution against analyzing data with low  $r^2$  values and so we considered only  $\delta^{18}\text{O}_{\text{R}}$  values with  $r^2 > 0.8$  (resulting in the elimination of 5 of the 22 observations). For both forest and pasture sites, we then calculated the  $\delta^{18}\text{O}_{\text{R}}$  values expected for a  $\text{CO}_2$  efflux in equilibrium with stem (soil) water (calculated using Eq. 6).

There was a weak correlation between the observed  $\delta^{18}\text{O}_{\text{R}}$  and that predicted for  $\text{CO}_2$  in equilibrium with soil water only (as predicted by Eq. 6), although the correlation was generally stronger for the wet season than the dry season (Fig. 9). By contrast there was a strong positive correlation between the observed  $\delta^{18}\text{O}_{\text{R}}$  values and the difference between observed  $\delta^{18}\text{O}_{\text{R}}$  and that predicted by Eq. 6 (Fig. 10). In general the differences between predicted and observed  $\delta^{18}\text{O}_{\text{R}}$  values were greater in the dry season at both forest and pasture sites. This observation suggests that leaf water remains enriched above that of soil (and stem water) at night in these tropical ecosystems and that leaf respiration




FIG. 7. Relationship between the oxygen isotope ratio of  $\text{CO}_2$  in air samples and  $1/[\text{CO}_2]$  ( $\text{ppm}^{-1}$ ) (Keeling plot) collected from the head space of a soil respiration chamber in the Santarém primary forest site in Brazil during March 2001.




FIG. 8. Comparison of the temporal variation in the oxygen isotope ratio of stem water and the isotope ratio of  $\text{CO}_2$  respired at night by the entire ecosystem in forest and pasture sites in Brazil. Data are means  $\pm$  1 SE.

contributes a large proportion of total ecosystem respired  $\text{CO}_2$ . In addition, the relative magnitudes of leaf and soil respiration may change between wet and dry seasons, so that the influence of leaf respiration on the oxygen isotope ratio of total ecosystem respiration increases during the dry season.

#### Modeling $\Delta\text{C}^{18}\text{OO}$ in forest and pasture ecosystems

Measurements of  $\delta^{18}\text{O}_{\text{Leaf}}$  values were used in Eqs. 4 and 5 to make calculations of the midday values for discrimination against  $\text{C}^{18}\text{OO}$  during photosynthetic gas exchange ( $\Delta\text{C}^{18}\text{OO}$ ). Forests had consistently higher modeled  $\Delta\text{C}^{18}\text{OO}$  values than pastures (Fig. 11). This resulted from similar  $\delta^{18}\text{O}_{\text{Leaf}}$  values, but lower  $c_c$  and much lower  $\theta$  values associated with reduced carbon anhydrase (CA) activity in  $\text{C}_4$  grasses (Gillon and Yakir 2001). The temporal variation in Fig. 11 was caused completely by changes in  $\delta^{18}\text{O}_{\text{Leaf}}$  values.

Modeled  $\Delta\text{C}^{18}\text{OO}$  values were also used to compare forest and pasture sites in Santarém during September 2000, based on the calculated variation in  $\delta^{18}\text{O}_{\text{Leaf}}$  values shown in Fig. 6. These data indicated that average midday  $\Delta\text{C}^{18}\text{OO}$  values were  $\sim 20\text{‰}$  in the forest, significantly higher than the  $\sim 9\text{‰}$  calculated for the  $\text{C}_4$  grasses of the pasture site (Fig. 12).

#### DISCUSSION

Our study represents the first long-term, integrated set of isotopic measurements of water and  $\text{CO}_2$  for tropical rain forest and pasture ecosystems. These data provide insight into the factors driving changes in the  $\delta^{18}\text{O}$  values of atmospheric  $\text{CO}_2$ .

We observed good agreement between measured and predicted  $\delta^{18}\text{O}_{\text{Leaf}}$  values in the Manaus forest canopy (Fig. 3). Consistent with the Craig-Gordon (1965) model predictions as modified by Flanagan et al. (1991),




FIG. 9. Comparison between the observed oxygen isotope ratio of  $\text{CO}_2$  respired at night by the entire ecosystem and the predicted isotope ratio of respired  $\text{CO}_2$  calculated using Eq. 6. The calculations done using Eq. 6 assume that all respired  $\text{CO}_2$  has equilibrated with water that has the same  $\delta^{18}\text{O}$  value as that measured for stem (soil) water at a site. Values are shown separately for the wet and dry seasons for all forest and pasture sites sampled in Brazil.




FIG. 10. Comparison of the difference between the oxygen isotope ratio ( $\delta^{18}\text{O}$ , ‰ SMOW) expected of respired  $\text{CO}_2$  if all water sources were in equilibrium with soil (stem) water  $\text{CO}_2$  and nighttime field observations as a function of the oxygen isotope ratio of ecosystem respiration. The calculations were performed using Eq. 6 and assume that all respired  $\text{CO}_2$  has fully equilibrated with water that has the same  $\delta^{18}\text{O}$  value as that measured for stem (soil) water at a site. Values represent all observations from forest and pasture sites during both the wet and dry seasons where  $r^2 > 0.8$ .

there were two main factors that contributed to variation in forest leaf water  $\delta^{18}\text{O}$  values, height variation within a forest canopy and wet vs. dry season differences. Both of these factors were primarily controlled by changes in vapor pressure deficit (VPD), although from one sample date to the next there were changes in the isotope composition of source water (Flanagan et al. 1991, Roden and Ehleringer 1999). Our leaf water  $\delta^{18}\text{O}$  observations agree with and extend previous shorter term observations for tropical forest regions. Zundel et al. (1978) reported diurnal  $\delta^{18}\text{O}_{\text{Leaf}}$  values for two trees in a tropical rain forest near Itabuna ( $14^\circ \text{S}$ ,

$39^\circ \text{W}$ ); maximum  $\delta^{18}\text{O}_{\text{Leaf}}$  values of 6–10‰ were observed (with a source water value of  $-2\text{‰}$ ). Interestingly, Zundel et al. (1978) also reported that  $\delta^{18}\text{O}_{\text{Leaf}}$  values from semiarid caatinga forests to the south reached  $+14\text{‰}$  at midday (with a source water value of  $-4\text{‰}$ ). Moreira et al. (1997) and Sternberg et al. (1998) reported  $\delta^{18}\text{O}_{\text{Leaf}}$  values from primary forest leaves at the Ducke Reserve near Manaus, Brazil, that were isotopically lighter than those observed in this study. In all likelihood, those differences may have been associated with lower VPD values during the sampling periods. Sternberg et al. (1989) also found significant height gradients in the  $\delta^{18}\text{O}$  of leaf cellulose at Barro Colorado Island, Panama. The height gradients in cellulose  $\delta^{18}\text{O}$  were associated with humidity gradients within the Panama tropical moist forest canopy.

In our study, leaf water  $^{18}\text{O}$  enrichment in pastures typically exceeded that of adjacent primary forests, largely because of a greater VPD in pasture than in forest sites. The  $\delta^{18}\text{O}$  values of source waters reflect precipitation inputs; these values in pastures and in adjacent forests were not different from each other and, therefore, it was primarily variation in leaf water enrichment between sites that contributed to the different leaf water  $\delta^{18}\text{O}$  values observed through time. In addition to environmental variation (VPD) between pasture and forest sites, leaf water isotopic composition in the  $\text{C}_4$  pasture grasses is affected by progressive enrichment along parallel veins (Helliker and Ehleringer 2000). Grass  $\delta^{18}\text{O}_{\text{Leaf}}$  values can often be significantly higher in  $^{18}\text{O}$  relative to predictions of the Craig-Gordon (1965) model due to this progressive enrichment. Because of the limited availability of meteorological data at the remote study sites during this investigation, we were unable to calculate predicted  $\delta^{18}\text{O}_{\text{Leaf}}$  values in pasture sites for most of our sampling trips. Thus, we were unable to document the extent of the deviation between Craig-Gordon (1965) model predictions and observed leaf water  $\delta^{18}\text{O}$  values under field conditions in the tropical pastures.




FIG. 11. Temporal variation in discrimination against  $\text{CO}_2$  molecules containing  $^{18}\text{O}$  ( $\Delta\text{C}^{18}\text{OO}$ , ‰ SMOW) at midday in forest and pasture sites in the Amazon Basin. These data represent calculations involving Eq. 4 and using measured midday values of  $\delta^{18}\text{O}$  of leaf water and the following values:  $c_c = 282 \mu\text{mol/mol}$  for  $\text{C}_3$  trees,  $c_c = 180 \mu\text{mol/mol}$  for  $\text{C}_4$  grasses, and  $c_a = 370 \mu\text{mol/mol}$ , where  $c$  is the  $\text{CO}_2$  concentration in the air (a) and sites of carboxylation (c). We assume an equilibration value of  $\theta = 0.93$  for  $\text{C}_3$  trees and shrubs and  $\theta = 0.3$  for  $\text{C}_4$  grasses.




FIG. 12. Comparison of modeled discrimination against  $\text{CO}_2$  molecules containing  $^{18}\text{O}$  ( $\Delta\text{C}^{18}\text{OO}$ , ‰) in the Santarém forest and pasture sites during September 2000. The left-hand panels represent the mean diurnal pattern for the 10-d period shown in the right-hand panels. The calculations were done using Eq. 4 and the modeled leaf water  $\delta^{18}\text{O}$  values shown in Fig. 6;  $c_c = 282 \mu\text{mol/mol}$  for  $\text{C}_3$  trees,  $c_c = 180 \mu\text{mol/mol}$  for  $\text{C}_4$  grasses, and  $c_a = 370 \mu\text{mol/mol}$ . We assume that  $\theta = 0.93$  for  $\text{C}_3$  trees and shrubs and  $\theta = 0.3$  for  $\text{C}_4$  grasses.

Additional errors can occur in the application of a steady-state model of leaf water evaporative enrichment under field conditions. During the day, the  $^{18}\text{O}$  content of leaf water may lag steady-state values by as little as 1–2 h (Flanagan et al. 1991, Roden and Ehleringer 1999), but that time lag is greatly increased at night when water flow through the leaf is substantially reduced (Cernusak et al. 2002). The assumption of isotopic steady state at night will result in underestimation of the  $\delta^{18}\text{O}_{\text{Leaf}}$  values and this has implications for interpreting Keeling plot estimates of the oxygen isotope ratio of ecosystem respired  $\text{CO}_2$  (Flanagan et al. 1997, 1999).

We compared our observed  $\delta^{18}\text{O}_{\text{R}}$  values with those predicted based on  $\text{CO}_2$  in equilibrium with soil (stem) water. The predicted  $\delta^{18}\text{O}_{\text{R}}$  values also assumed a fractionation factor (8.8‰) during diffusion of  $\text{CO}_2$  (Tans 1998). A number of factors will contribute to variation between Keeling plot estimates and simple predictions of the oxygen isotope ratio of respired  $\text{CO}_2$ . For example, Tans (1998) has shown that invasion flux of atmospheric  $\text{CO}_2$  into the soil can influence the equilibration between soil  $\text{CO}_2$  and water, a result that causes curvature in the  $\delta^{18}\text{O}$  and  $1/[\text{CO}_2]$  relationship (at very high  $[\text{CO}_2]$ ). Therefore, the linear regression technique (Keeling plot) should result in  $\delta^{18}\text{O}_{\text{R}}$  values that are slightly more negative than actual  $\delta^{18}\text{O}_{\text{R}}$  values (Tans 1998). Application of a lower value (7.3‰ rather than 8.8‰) for the effective fractionation during diffusion of  $\text{CO}_2$  out of the soil can help deal with this problem (Miller et al. 1999, Stern et al. 1999, Yakir and Sternberg 2000). Leaf respiration at night will also affect

the oxygen isotope ratio of total ecosystem respired carbon dioxide, and leaf water  $\delta^{18}\text{O}$  values can be significantly different than those of the soil water. The Flanagan et al. (1991) leaf water model predicted lower  $\delta^{18}\text{O}_{\text{Leaf}}$  values at night, which were often not consistent with observed  $\delta^{18}\text{O}_{\text{R}}$  values. The difference may be explained by non steady-state conditions and the time lags necessary for the daytime  $^{18}\text{O}$  enrichment signal to be “washed” out of the leaf by input of unenriched xylem water (Cernusak et al. 2002).

During the dry season, aboveground vegetation appeared to contribute more to the oxygen isotope ratio of ecosystem respiration than soil respiration. We base this conclusion on Keeling plot intercept observations, assuming that  $\text{CO}_2$  fluxes were derived primarily from soil and leaf sources (Figs. 8, 9). Leaf water isotopic enrichment was generally greater in the dry season than in the wet season. In this case  $\delta^{18}\text{O}_{\text{Leaf}}$  values may also remain more enriched at night and affect the Keeling plot  $\delta^{18}\text{O}_{\text{R}}$  values to a greater extent in the dry season. Alternatively, there may be wet vs. dry season changes in the relative magnitude of leaf and soil respiration in these tropical forests. Preliminary eddy covariance and chamber data from the Santarém primary forest suggests that soil respiration is significantly reduced during the dry season and this contributes to higher net ecosystem  $\text{CO}_2$  uptake because of only minor seasonal variation in gross photosynthesis (Saleska et al. 2003, da Rocha et al. 2004).

Cernusak et al. (2002) provided strong experimental evidence that nighttime  $\delta^{18}\text{O}_{\text{Leaf}}$  values did not ever reach the expected steady-state values with their en-

vironment, most likely because the time to isotopic equilibration lengthened when stomata closed at night. The observed–modeled differences shown in Fig. 10 are consistent with the calculation not accounting for an  $^{18}\text{O}$  enriched source of respiratory  $\text{CO}_2$  in the forest and pasture ecosystems. Additionally, the increased deviations in expected and observed  $\delta^{18}\text{O}_R$  values with increasing  $\delta^{18}\text{O}_R$  values are again consistent with not accounting for  $\text{CO}_2$  efflux from enriched  $\delta^{18}\text{O}_{\text{Leaf}}$  values. This is because we would expect that when daytime humidity was low the  $^{18}\text{O}$  enrichment of leaf water above soil water would be greater and result in an even larger deviation from assumed steady-state values at night (Eq. 2). However, as no nighttime  $\delta^{18}\text{O}_{\text{Leaf}}$  values were available, we are unable to fully resolve this issue. It is also likely that the relative magnitudes of leaf and soil respiration may change between the wet and dry seasons, such that the influence of leaf respiration on the oxygen isotope ratio of total ecosystem respiration increases during the dry season in the forest sites. Saleska et al. (2003) and da Rocha et al. (2004) and provide evidence indicating that the contribution of soil  $\text{CO}_2$  efflux to the total ecosystem respiratory flux declines during the dry season. Thus, it seems inappropriate to conclude that the soil water  $^{18}\text{O}$  value alone dominates the  $\delta^{18}\text{O}_R$  values in these forest and pasture ecosystems, but rather it is likely that the seasonal variations are influenced by both aboveground and soil components as well as by possible changes in the partitioning of the relative fractions of  $\text{CO}_2$  derived from these distinct sources. Our focus here was not to evaluate different predictions from the multiple models now available in the literature to predict the  $\delta^{18}\text{O}$  values of soil  $\text{CO}_2$  efflux. Instead our intent was to point out that a simple assumed equilibrium calculation was unable to capture much of the observed seasonal isotopic variation.

Forests had consistently higher modeled  $\Delta\text{C}^{18}\text{OO}$  values than pastures. This resulted from similar  $\delta^{18}\text{O}_{\text{Leaf}}$  values, but lower  $c_c$  and much lower  $\theta$  values associated with reduced carbon anhydrase (CA) activity in  $\text{C}_4$  grasses (Gillon and Yakir 2001). With an increasing extent of forest to pasture conversion occurring in the Amazon Basin, it is possible that the reduced  $\Delta\text{C}^{18}\text{OO}$  associated with pasture vegetation could contribute to the decreasing global trend in the  $\delta^{18}\text{O}$  value of atmospheric  $\text{CO}_2$  as described by Gillon and Yakir (2001).

Model calculations by Ciais et al. (1997a, b) suggested a strong depletion of  $^{18}\text{O}$  in atmospheric  $\text{CO}_2$  for tropical regions covered by evergreen vegetation, in both Africa and South America (Amazon Basin). The  $\delta^{18}\text{O}$  values of atmospheric  $\text{CO}_2$  predicted for the continental tropical regions with high productivity were even lower than the values calculated for the very high northern latitudes. This suggests that the oxygen isotope ratio of ecosystem respired  $\text{CO}_2$  must be significantly depleted in  $^{18}\text{O}$  compared to the enrichment

of  $^{18}\text{O}$  in  $\text{CO}_2$  caused by discrimination during photosynthesis in these regions. Our model calculations, which utilize observed leaf water values, predict that  $\Delta\text{C}^{18}\text{OO}$  values for pasture and forest should vary between  $\sim 8\text{--}20\%$  (Fig. 12). These discrimination values are slightly lower than the average  $\Delta\text{C}^{18}\text{OO}$  values of  $\sim 21\%$  observed during the growing season months in the boreal forest regions of Canada (Flanagan et al. 1997).

Low VPD values generally limit the extent of leaf water isotopic enrichment in tropical forests compared to boreal regions and so  $\Delta\text{C}^{18}\text{OO}$  values are low despite the high leaf  $c_c$  values indicated by leaf  $\delta^{13}\text{C}$  values (Ometto et al. 2002). In addition, the average forest  $\delta^{18}\text{O}_R$  value observed in this study was  $\sim 28\%$  on the SMOW scale (Fig. 8), which is equivalent to a  $\delta^{18}\text{O}_R$  of  $-13\%$  on the PDB- $\text{CO}_2$  scale. This is quite close to the average  $\delta^{18}\text{O}_R$  value of  $-14.1\%$  (PDB- $\text{CO}_2$ ) determined by Flanagan et al. (1997) for Canadian boreal forest vegetation. Given the similar values for the isotope effects during photosynthesis and respiration in tropical and boreal regions, and the much higher  $\text{CO}_2$  fluxes in tropical evergreen forests, it is expected that atmospheric  $\text{CO}_2$  should be depleted in  $^{18}\text{O}$  over the Amazon Basin region as predicted by Ciais et al. (1997a, b). Aircraft sampling of the  $\delta^{18}\text{O}$  values of atmospheric  $\text{CO}_2$  over the Amazon Basin is required to further test these ideas.

Long-term, integrated sets of isotopic measurements of water and  $\text{CO}_2$  from terrestrial ecosystems are essential for constraining our interpretations of global and regional carbon cycles. Here we have presented the first multiseasonal data sets for tropical rain forest and pasture ecosystems. These data sets are consistent with predicted patterns proposed by Ciais et al. (1997a, b) and provide insight into the biological factors driving changes in the  $\delta^{18}\text{O}$  values of atmospheric  $\text{CO}_2$ .

#### ACKNOWLEDGMENTS

This study has been supported by a grant from NASA LBA-Ecology as part of the Large Scale Biosphere Atmosphere Experiment in Amazonia. We thank the personnel at CENA-USP for assistance with stable isotope ratio analyses. We thank Michael Goulden, Scott Miller, Ricardo Sakai, David Fitzgarrald, and Ricardo Della Rosa for meteorological data.

#### LITERATURE CITED

- Behrenfeld, M. J., J. T. Randerson, C. R. McClain, G. C. Feldman, S. O. Los, C. J. Tucker, P. G. Falkowski, C. B. Field, R. Frouin, W. E. Esaias, and D. D. Kolber. 2001. Biospheric primary production during an ENSO transition. *Science* **291**:2594–2597.
- Bowling, D. R., C. S. Cook, and J. R. Ehleringer. 2001. Technique to measure  $\text{CO}_2$  mixing ratio in small flasks with a bellows/IRGA system. *Agricultural and Forest Meteorology* **109**:61–65.
- Bowling, D. R., N. G. McDowell, B. V. Bond, B. E. Law, and J. R. Ehleringer. 2002.  $^{13}\text{C}$  content of ecosystem respiration is linked to precipitation and vapor pressure deficit. *Oecologia* **131**:113–124.
- Bowling, D. R., N. G. McDowell, J. M. Welker, B. J. Bond, B. E. Law, and J. R. Ehleringer. 2003. Oxygen isotope content of  $\text{CO}_2$  in nocturnal ecosystem respiration: 2. Short-

- term dynamics of foliar and soil component fluxes in an old-growth ponderosa pine forest. *Global Biogeochemical Cycles* **17**(4):1124. [doi:10.1029/2003GB002082]
- Buchmann, N., J.-M. Guehl, T. S. Barigah, and J. R. Ehleringer. 1997. Interseasonal comparison of CO<sub>2</sub> concentrations, isotopic composition, and carbon dynamics in an Amazonian rainforest (French Guiana). *Oecologia* **110**: 120–131.
- Canadell, J., et al. 2000. Carbon metabolism of the terrestrial biosphere: a multitechnique approach for improved understanding. *Ecosystems* **3**:115–130.
- Cernusak, L. A., J. S. Pate, and G. D. Farquhar. 2002. Diurnal variation in the stable isotope composition of water and dry matter in fruiting *Lupinus angustifolius* under field conditions. *Plant Cell and Environment* **25**:893–907.
- Ciais, P., et al. 1997a. A three-dimensional synthesis study of δ<sup>18</sup>O in atmospheric CO<sub>2</sub>. 1. Surface fluxes. *Journal of Geophysical Research* **102**:5857–5872.
- Ciais, P., et al. 1997b. A three-dimensional synthesis study of δ<sup>18</sup>O in atmospheric CO<sub>2</sub>. 2. Simulations with the TM2 transport model. *Journal of Geophysical Research* **102**: 5873–5883.
- Ciais, P., and H. A. J. Meijer. 1998. The <sup>18</sup>O/<sup>16</sup>O isotope ratio of atmospheric CO<sub>2</sub> and its role in global cycle research. Pages 409–431 in H. Griffiths, editor. *Stable isotopes: integration of biological, ecological and geochemical research*. BIOS Scientific Publishers, Oxford, UK.
- Craig, H., and L. I. Gordon. 1965. Deuterium and oxygen-18 variations in the ocean and the marine atmosphere. Pages 9–130 in E. Tongiogi, editor. *Proceedings of a conference on stable isotopes in oceanographic studies and paleotemperatures*. V. Lischi and Figli, Pisa, Italy.
- Cramer, W., D. W. Kicklighter, A. Bondeau, B. Moore, III, G. Churkina, B. Nemry, A. Ruimy, A. L. Schloss, and the Participants of the Potsdam NPP Model Intercomparison. 1999. Comparing global models of terrestrial net primary productivity (NPP): overview and key results. *Global Change Biology* **5**:1–15.
- Cramer, W., et al. 2001. Global response of terrestrial ecosystem structure and function to CO<sub>2</sub> and climate change: results from six dynamic global vegetation models. *Global Change Biology* **7**:357–373.
- da Rocha, H. R., M. L. Goulden, S. D. Miller, M. C. Menton, L. D. V. O. Pinto, H. C. Freitas, and A. M. S. Figueira. 2004. Seasonality of water and heat fluxes over a tropical forest in eastern Amazonia. *Ecological Applications* **14**: S22–S32.
- Ehleringer, J. R., D. R. Bowling, L. B. Flanagan, J. Fessenden, B. Helliker, L. A. Martinelli, and J. P. Ometto. 2002. Stable isotopes and carbon cycle processes in forests and grasslands. *Plant Biology* **4**:181–189.
- Ehleringer, J. R., and C. S. Cook. 1998. Carbon and oxygen isotope ratios of ecosystem respiration along an Oregon conifer transect: preliminary observations based on small-flask sampling. *Tree Physiology* **18**:513–519.
- Ehleringer, J. R., J. Roden, and T. E. Dawson. 2001. Assessing ecosystem-level water relations through stable isotope ratio analyses. Pages 181–198 in O. E. Sala, R. B. Jackson, H. A. Mooney, and R. W. Howarth, editors. *Methods in ecosystem science*. Springer Verlag, New York, New York, USA.
- Farquhar, G. D., J. R. Ehleringer, and K. T. Hubick. 1989. Carbon isotope discrimination and photosynthesis. *Annual Review of Plant Physiology and Molecular Biology* **40**: 503–537.
- Farquhar, G. D., and J. Lloyd. 1993. Carbon and oxygen isotope effects in the exchange of carbon dioxide between plants and the atmosphere. Pages 47–70 in J. R. Ehleringer, A. E. Hall, and G. D. Farquhar, editors. *Stable isotopes, and plant carbon–water relations*. Academic Press, New York, New York, USA.
- Farquhar, G. D., J. Lloyd, J. A. Taylor, L. B. Flanagan, J. P. Syvertsen, K. T. Hubick, S. C. Wong, and J. R. Ehleringer. 1993. Vegetation effects on the isotope composition of oxygen in atmospheric CO<sub>2</sub>. *Nature* **363**:439–443.
- Fessenden, J. E., C. S. Cook, and J. R. Ehleringer. 2002. Rapid <sup>18</sup>O analysis of small sized water samples using a continuous-flow isotope ratio mass spectrometer. *Rapid Communications in Mass Spectrometry* **16**:1257–1260.
- Fessenden, J. E., and J. R. Ehleringer. 2002. Age-related variations in δ<sup>13</sup>C of ecosystem respiration across a coniferous forest chronosequence in the Pacific Northwest. *Tree Physiology* **22**:159–167.
- Flanagan, L. B., J. R. Brooks, G. T. Varney, and J. R. Ehleringer. 1997. Discrimination against C<sup>18</sup>O<sup>16</sup>O during photosynthesis and the oxygen isotope ratio of respired CO<sub>2</sub> in boreal forest ecosystems. *Global Biogeochemical Cycles* **11**:83–98.
- Flanagan, L. B., J. P. Comstock, and J. R. Ehleringer. 1991. Comparison of modeled and observed environmental influences on the stable oxygen and hydrogen isotope composition of leaf water in *Phaseolus vulgaris* L. *Plant Physiology* **96**:588–596.
- Flanagan, L. B., and J. R. Ehleringer. 1998. Ecosystem-atmosphere CO<sub>2</sub> exchange: interpreting signals of change using stable isotope ratios. *Trends in Ecology and Evolution* **13**:10–14.
- Flanagan, L. B., D. S. Kubien, and J. R. Ehleringer. 1999. Spatial and temporal variation in the carbon and oxygen stable isotope ratio of respired CO<sub>2</sub> in a boreal forest ecosystem. *Tellus* **51B**:367–384.
- Francey, R. J., and P. P. Tans. 1987. Latitudinal variation in oxygen-18 of atmospheric CO<sub>2</sub>. *Nature* **327**:495–497.
- Friedli, H., U. Siegenthaler, D. Rauber, and H. Oeschger. 1987. Measurements of concentration, <sup>13</sup>C/<sup>12</sup>C and <sup>18</sup>O/<sup>16</sup>O ratios of tropospheric carbon dioxide over Switzerland. *Tellus* **39**:80–88.
- Fung, I. 2000. Climate change: variable carbon sinks. *Science* **290**:1313.
- Gillon, J., and D. Yakir. 2001. Influence of carbonic anhydrase activity in terrestrial vegetation on the <sup>18</sup>O content of atmospheric CO<sub>2</sub>. *Science* **291**:2584–2587.
- Helliker, B. R., and J. R. Ehleringer. 2000. Establishing a grassland signature in veins: <sup>18</sup>O in the leaf water of C<sub>3</sub> and C<sub>4</sub> grasses. *Proceedings of the National Academy of Sciences* **97**:7894–7898.
- Helliker, B. R., J. R. Roden, C. Cook, and J. R. Ehleringer. 2002. A rapid and precise method for sampling and determining the oxygen isotope ratio of atmospheric water vapor. *Rapid Communications in Mass Spectrometry* **16**: 929–932.
- Hodnett, M. G., M. D. Oyama, J. Tomazella, and A. de O. Marques Filho. 1996. Comparisons of long term soil water storage behavior under pasture and forest in three areas of Amazonia. Pages 79–100 in J. H. C. Gash, C. A. Nobre, J. M. Roberts, and R. L. Victoria, editors. *Amazonian deforestation and climate*. John Wiley and Sons, Chichester, UK.
- Keeling, C. D. 1958. The concentration and isotopic abundances of atmospheric carbon dioxide in rural areas. *Geochimica et Cosmochimica Acta* **13**:322–334.
- Majoube, M. 1971. Fractionnement en oxygene 18 et en deuterium entre l'eau et sa vapeur. *Journal de Chimie et Physique* **58**:1423–1436.
- Marland, G., A. Brenkert, and J. Olivier. 1999. CO<sub>2</sub> from fossil fuel burning: a comparison of ORNL and EDGAR estimates of national emissions. *Environmental Science and Policy* **2**:265–273.

- Miller, J. B., D. Yakir, J. W. C. White, and P. P. Tans. 1999. Measurement of  $^{18}\text{O}/^{16}\text{O}$  in the soil-atmosphere  $\text{CO}_2$  flux. *Global Biogeochemical Cycles* **13**:761–774.
- Moreira, M. Z., L. S. L. Sternberg, L. A. Martinelli, R. L. Victoria, E. M. Barbosa, C. M. Bonates, and D. C. Nepstad. 1997. Contribution of transpiration to forest ambient vapor based on isotopic measurements. *Global Change Biology* **3**:439–450.
- Ometto, J. P. H. B., L. B. Flanagan, L. A. Martinelli, M. Z. Moreira, N. Higuchi, and J. R. Ehleringer. 2002. Carbon isotope discrimination in forest and pasture ecosystems of the Amazon Basin, Brazil. *Global Biogeochemical Cycles* **16**(4):1109. [doi:10.1029/2001GB001462]
- Pataki, D. E., J. R. Ehleringer, L. B. Flanagan, D. Yakir, D. R. Bowling, C. J. Still, N. Buchmann, J. O. Kaplan, and J. A. Berry. 2003. The application and interpretation of Keeling plots in terrestrial carbon cycle research. *Global Biogeochemical Cycles* **17**:1022. [doi:10.1029/2001GB001850]
- Peylin, P., P. Ciais, A. S. Denning, P. P. Tans, J. A. Berry, and J. W. C. White. 1999. A 3-dimensional study of the  $\delta^{18}\text{O}$  in atmospheric  $\text{CO}_2$ : contribution of different land ecosystems. *Tellus* **51B**:642–667.
- Riley, W. J., C. J. Still, M. S. Torn, and J. A. Berry. 2002. A mechanistic model of  $\text{H}_2^{18}\text{O}$  and  $\text{C}^{18}\text{OO}$  fluxes between ecosystems and the atmosphere: model description and sensitivity analyses. *Global Biogeochemical Cycles* **16**(4):1095. [doi:1029/2002GB001878]
- Roden, J. S., and J. R. Ehleringer. 1999. Observations of hydrogen and oxygen isotopes in leaf water confirm the Craig-Gordon model under wide-ranging environmental conditions. *Plant Physiology* **120**:1165–1173.
- Roden, J. S., G. Lin, and J. R. Ehleringer. 2000. A mechanistic model for interpretation of hydrogen and oxygen isotope ratios in tree-ring cellulose. *Geochimica et Cosmochimica Acta* **64**:21–35.
- Saleska, S. R., et al. 2003. Carbon in Amazon forests: unexpected seasonal fluxes and disturbance-induced losses. *Science* **302**:1554–1557.
- Socki, R. A., C. S. Romanek, and E. K. Gibson. 1999. On-line technique for measuring stable oxygen and hydrogen isotopes from microliter quantities of water. *Analytical Chemistry* **71**:2250–2253.
- Stern, L., W. T. Baisden, and R. Amundson. 1999. Processes controlling the oxygen isotope ratio of soil  $\text{CO}_2$ : analytic and numeric modeling. *Geochimica et Cosmochimica Acta* **63**:799–814.
- Sternberg, L. S. L., M. Z. Moreira, L. A. Martinelli, R. L. Victoria, E. M. Barbosa, L. C. M. Bonates, and D. C. Nepstad. 1998. The relationship between  $^{18}\text{O}/^{16}\text{O}$  and  $^{13}\text{C}/^{12}\text{C}$  ratios of ambient  $\text{CO}_2$  in two Amazonian tropical forests. *Tellus* **50B**:366–376.
- Sternberg, L. S. L., S. S. Mulkey, and S. J. Wright. 1989. Oxygen isotope ratio stratification in a tropical moist forest. *Oecologia* **81**:51–56.
- Tans, P. P. 1998. Oxygen isotopic equilibrium between carbon dioxide and water in soils. *Tellus* **50B**:163–178.
- Tans, P. P., and J. W. C. White. 1998. In balance, with a little help from the plants. *Science* **281**:183–184.
- Tian, H., J. M. Melillo, D. W. Kicklighter, A. D. McGuire, J. V. K. Helfrich, B. Moore, and C. J. Vorosmarty. 1998. Effect of interannual climate variability on carbon storage in Amazonian ecosystems. *Nature* **396**:664–667.
- Yakir, D., and L. S. L. Sternberg. 2000. The use of stable isotopes to study ecosystem gas exchange. *Oecologia* **123**:297–311.
- Zundel, G., W. Miekeley, B. M. Grisi, and H. Forstel. 1978. The  $\text{H}_2^{18}\text{O}$  enrichment in the leaf water of tropic trees: comparison of species from the tropical rain forest and the semi-arid region in Brazil. *Radiation and Environmental Biophysics* **15**:203–212.

#### SUPPLEMENT

Files of data collected at, or relevant to, different field sites that were used to construct the Keeling plot intercept figures and  $^{18}\text{O}$  discrimination figures published in the text are available in ESA's Electronic Data Archive: *Ecological Archives* A015-003-S1.