

J. Canadell · R.B. Jackson · J.R. Ehleringer  
H.A. Mooney · O.E. Sala · E.-D. Schulze

## Maximum rooting depth of vegetation types at the global scale

Received: 26 January 1996 / Accepted: 18 July 1996

**Abstract** The depth at which plants are able to grow roots has important implications for the whole ecosystem hydrological balance, as well as for carbon and nutrient cycling. Here we summarize what we know about the maximum rooting depth of species belonging to the major terrestrial biomes. We found 290 observations of maximum rooting depth in the literature which covered 253 woody and herbaceous species. Maximum rooting depth ranged from 0.3 m for some tundra species to 68 m for *Boscia albitrunca* in the central Kalahari; 194 species had roots at least 2 m deep, 50 species had roots at a depth of 5 m or more, and 22 species had roots as deep as 10 m or more. The average for the globe was  $4.6 \pm 0.5$  m. Maximum rooting depth by biome was  $2.0 \pm 0.3$  m for boreal forest,  $2.1 \pm 0.2$  m for cropland,  $9.5 \pm 2.4$  m for desert,  $5.2 \pm 0.8$  m for sclerophyllous shrubland and forest,  $3.9 \pm 0.4$  m for temperate coniferous forest,  $2.9 \pm 0.2$  m for temperate deciduous forest,  $2.6 \pm 0.2$  m for temperate grassland,  $3.7 \pm 0.5$  m for tropical deciduous forest,  $7.3 \pm 2.8$  m for tropical evergreen forest,  $15.0 \pm 5.4$  m for tropical grassland/savanna, and  $0.5 \pm 0.1$  m for tundra. Grouping all the species across biomes (except croplands) by three basic functional groups: trees, shrubs, and herbaceous plants, the maximum rooting depth was  $7.0 \pm 1.2$  m

for trees,  $5.1 \pm 0.8$  m for shrubs, and  $2.6 \pm 0.1$  m for herbaceous plants. These data show that deep root habits are quite common in woody and herbaceous species across most of the terrestrial biomes, far deeper than the traditional view has held up to now. This finding has important implications for a better understanding of ecosystem function and its application in developing ecosystem models.

**Key words** Deep roots function · Terrestrial vegetation · Biomes · Plant forms · Root depth

### Introduction

There is good evidence that some plant species are able to send roots very deep in the soil. This pattern is indicated by plants that grow well into the summer drought and by desert plants that grow for years with minimal or no rainfall (Batanouny and Abdel Wahab 1973; Poole and Miller 1975). In fact, survivorship of some species in arid systems has been shown to depend completely on a plant's ability to tap water from permanent water tables, which are sometimes located at depths of 18 m or more (Ravitscher 1948; Lewis and Burghy 1964). In addition, there have been direct observations of roots at depths below 2–3 m in caves, road cuts, mine shafts and trenches, and in some instances, roots of woody species have been seen exceptionally deep in the soil. This is the case of *Boscia albitrunca* and *Acacia erioloba* whose roots have been found at a depth of 68 m and 60 m, respectively, in the central Kalahari, Botswana (Jennings 1974), and the case of mesquite roots (*Prosopis juliflora*) found at 53 m deep in the Sonoran Desert, United States (Phillips 1963). Similarly, Stone and Kalisz (1991) reported 11 tree species rooted below 20 m depth. Hence, we know of the potential of some species to have very deep roots at few sites, yet very little is known about how common the habit of deep rooting is across species and environments.

There are two main reasons why this below-ground aspect of ecosystem structure, with its important functional implications, has been under-emphasized. First of all, there are a number of studies on root biomass distri-

J. Canadell (✉) · H.A. Mooney  
Department of Biological Sciences, Stanford University,  
Stanford, CA 94305, USA  
fax: (415) 723-9253; email: jcanadel@leland.stanford.edu

R.B. Jackson  
Department of Botany, University of Texas at Austin,  
Austin, TX 78713, USA

J.R. Ehleringer  
Department of Biology, University of Utah,  
Salt Lake City, UT 84112, USA

O.E. Sala  
Departamento de Ecología, Facultad de Agronomía,  
Universidad de Buenos Aires, Av. San Martín 4453,  
Buenos Aires, Argentina

E.-D. Schulze  
Lehrstuhl Pflanzenökologie, Universität Bayreuth,  
Postfach 101251, D-954440 Bayreuth, Germany

bution that show that most of the root biomass occurs within the first 50 cm of the soil, and that only a minimal fraction reaches depths below that depth (for a recent review see Jackson et al. 1996). Therefore, it has been assumed that a good understanding of the role of the root system regarding structure and function at the ecosystem level can be achieved by studying only the first 0.5 m of soil. Secondly, after a whole century of research on root systems, the means of obtaining data on root distribution and structure has not changed substantially: methods include the manual digging of trenches, the use of various mechanical excavation devices, dynamite, or high pressure water. When it comes to looking at patterns of maximum rooting depth, some of that technology is not even sufficient to provide access to deeper soil layers.

The functional significance of deep roots and their contribution to whole-ecosystem processes is still poorly understood. However, there is an increasing body of research in this field that shows the major role of deep roots, particularly for ecosystem water fluxes, as well as for carbon and nutrient cycling (Nepstad et al. 1994; Fisher et al. 1994; Richter and Markewitz 1985; Trumbore et al. 1995; Dawson 1996; Schulze et al. 1996).

The main objective of this review is to summarize what we know about the maximum rooting depth of the major terrestrial biomes ranging from tundra to tropical forest. The data set presented here provides information on plant structure which is relevant for a better mechanistic understanding of ecosystem function.

## Methods

We selected references which had species- or community-level information on root depth below 1.0 m, except for the tundra biome for which we considered all depths because permafrost usually limits root growth beyond 30–50 cm. Here we included references from journal papers, books, reports, and unpublished data when relevant, which cover all continents except Antarctica. The major biomes we considered were: boreal forest, croplands, desert, sclerophyllous shrubland and forest, temperate coniferous forest, temperate deciduous forest, temperate grassland, tropical deciduous forest, tropical evergreen forest, tropical grassland/savanna, and tundra. The species were grouped by biome which means that in some instances two different functional groups, such as grasses and shrubs, may be in the same biome category. This was the case of the tropical grasslands and savannas where both herbaceous and woody species occur together. Similarly, in the temperate grassland we also found a few common shrub species along with the bulk of herbaceous plants. Finally, root data for the commonest agricultural crops were collected, including wheat, soybean, alfalfa, barley, and a few other species.

For each rooting depth observation, we recorded the species from which the observation was made, and the community's dominant species when roots were not identified at the species level. For most of the references, the maximum root depth observed corresponded with the depth of the trench, road cut, mine pit, or other excavation, and it is safe to say that roots probably reached much deeper layers than those recorded. Almost all the data presented here came from direct observations of roots in road cuts, mine shafts, open-cut mines and trenches, and only a few values were inferred from the results of isotopic trace studies or plant and soil water potential measurements. Finally, we also recorded the soil type or any soil textural attribute available to characterize the soil environment in which roots were growing.

## Results and discussion

### Maximum rooting depth across biomes


We compiled a total of 290 observations of rooting depth which covered 253 different plant species from 11 biomes around the world. From this data set, 194 species had roots at least 2 m deep, 50 species had roots at a depth of 5 m or more, and 22 species had roots as deep as 10 m or more (Appendix 1). The average maximum rooting depth for the globe was  $4.6 \pm 0.5$  m, and the individual maximum rooting depth was 68 m for *Boscia albitrunca*, the roots of which were found during well drilling in deep sandy soils in the central Kalahari, Botswana (Jennings 1974). The ten deepest rooting species were in decreasing order: *Boscia albitrunca* (68 m), *Acacia erioloba* (60 m), *Prosopis juliflora* (53 m), *Eucalyptus marginata* (40 m), *Retama raetam* (20 m), *Tamarix aphylla* (20 m), *Andira humilis* (18 m), *Alhagi maurorum* (15 m), *Prosopis farcta* (15 m), and *Prosopis glandulosa* (15 m).

Figure 1 shows the maximum rooting depth for all species across biomes in which only the deepest rooting depth is plotted when a given species has more than one observation. Maximum rooting depth by biome was  $2.0 \pm 0.3$  m ( $n = 6$ ; highest value = 3.3 m) for boreal forest,  $2.1 \pm 0.2$  m ( $n = 17$ ; highest value = 3.7 m) for cropland,  $9.5 \pm 2.4$  m ( $n = 22$ ; highest value = 53 m) for desert,  $5.2 \pm 0.8$  m ( $n = 57$ ; highest value = 40) for sclerophyllous shrubland and forest,  $3.9 \pm 0.4$  m ( $n = 17$ ; highest value = 7.5 m) for temperate coniferous forest,  $2.9 \pm 0.2$  m ( $n = 19$ ; highest value = 4.4 m) for temperate deciduous forest,  $2.6 \pm 0.2$  m ( $n = 82$ ; highest value = 6.3 m) for temperate grassland,  $3.7 \pm 0.5$  m ( $n = 5$ ; highest value = 4.7 m) for tropical deciduous forest,  $7.3 \pm 2.8$  m ( $n = 5$ ; highest value = 18 m) for tropical evergreen forest,  $15.0 \pm 5.4$  m ( $n = 15$ ; highest value = 68 m) for tropical grassland/savanna, and  $0.5 \pm 0.1$  m ( $n = 8$ ; highest value = 0.9 m) for tundra.


Grouping all the species across biomes (except croplands) by three basic functional groups: trees, shrubs, and herbaceous plants, the maximum rooting depth was  $7.0 \pm 1.2$  m ( $n = 82$ ) for trees,  $5.1 \pm 0.8$  m ( $n = 69$ ) for shrubs, and  $2.6 \pm 0.1$  m ( $n = 85$ ) for herbaceous plants (Fig. 2).

Although differences are large among biomes, there are also important departures from the mean rooting depth pattern within a biome. In the boreal forest, for instance, the water table usually limits the downward growth of roots of *Larix laricina* and *Picea mariana*, whose roots are commonly found no deeper than 0.3 m. Other species, however, do have the capacity to grow below the water table down to a depth of 2 m (Strong and La Roi 1983).

Plants from arid environments or from environments with a long dry season showed the deepest rooting habits of all. The presence of water at deep layers makes it possible for some plants to survive in the rainshadow environments by tapping water from layers as deep as 53 m in the desert of the southwestern United States (Phillips



**Fig. 1** Reported species maximum rooting depth (m) grouped by terrestrial biome. When there are more than one observations for a given species, only the maximum value is plotted



**Fig. 2** Mean and SE of reported maximum rooting depth (m) by three major functional groups (trees, shrubs, and herbaceous plants) and crops

1963), and from 68 m deep, possibly even from 140 m deep where the water table was located, in the dry savanna of the central Kalahari (Jennings 1974). Likewise a group of species which also has a consistent pattern of deep rooting is that of the sclerophyllous trees, mostly made up of *Eucalyptus* spp. and *Quercus* spp. from the various Mediterranean regions of the world. The mean maximum rooting depth for sclerophyllous trees is  $12.6 \pm 3.4$  m ( $n = 11$ ), with *E. marginata* in Australia the deepest of all at about 40 m (Dell et al. 1983). Sclerophyllous shrubs, although with deep rooting habits, has a shallower rooting pattern with a maximum rooting depth of  $3.5 \pm 0.3$  m ( $n = 48$ ).

It is generally thought that roots in the evergreen tropical forest tend to be very shallow, but in this review the mean maximum rooting depth of 6 observations is  $6.5 \pm 2.5$  m. The only study that presented data from a depth beyond a few meters in the tropical forests of Brazil found roots all the way down to 18 m deep (Nepstad et al. 1994).

Another surprising result is the depth to which roots of herbaceous plants can descend, that was  $2.4 \pm 0.1$  m as average in this review. Weaver (1919) has published the most complete study to date, on rooting depth habits of herbaceous plants in a prairie in Nebraska, United States. Of 33 species he studied, 18 species have roots that extend beyond depths of 1.5 m, most of them between 2.1 m and 2.7 m, and few to a maximum depth of from 4 m to 6.1 m.

These results offer plenty of evidence that many plant species have the capacity for deep rooting in the soil, and

they provide enough data to challenge the dogma that plants are shallow rooted. Here we have presented, however, data on maximum rooting depth for individuals with the greatest depth. This value represents the observed maximum capacity of a given species to send roots deep into the soil, depths which may be reached by a small number of species and/or individuals within a community. In addition, an average or community weighted maximum rooting depth would also be functionally significant, yet data regarding this are hardly available for any biome. To illustrate the differences between absolute and average maximum rooting depth we shall present data from the root atlas published by Kutschera (1960). A random selection of 69 dicotyledonous species from grasslands in Mid Europe have an average maximum rooting depth of 1.1 m, the average of the 10 deepest species is 4.2 m and the absolute maximum rooting depth is 6.3 m. The average maximum rooting depth, which is the measurement most relevant to ecosystem functioning, will depend on species composition and density, and soil characteristics, all of which are fairly variable in space.

### Getting very deep

Plants show a variety of root types through which they have access to deep soil layers. The most common are tap roots, sinker roots and obliquely descending lateral roots, all of them important adaptations for reaching deep soils. The phenotypic expression of these root types is species dependent, but environmental conditions may completely change root structure, architecture, and depth to which roots are able to descend (Feldman 1984). Tap roots are probably the most specialized root type to access and transport water from deep soil horizons. Tap roots are very common across species and they were found in up to 75% of tropical trees (Klinge 1973), in 73 of 100 Mediterranean woody species (Canadell and Zedler 1995), and in 19 of 30 herbaceous species in the Rocky Mountains foothills, United States (Holch et al. 1941).

The downward growth of roots can be limited by a variety of factors, such as soil bulk density or shallow bedrock, but probably the most efficient barriers are horizontally stratified layers of shale or clay, permafrost, and water table (Dennis et al. 1978; Bennie 1991). There is a common notion that deep roots are mainly limited to sandy loose soils where mechanical impedance to root penetration is least. On the contrary, we have reported in this review a number of examples in which plants have found their way down to very deep layers, even in compact clay and rocky soils, and through hard pans (Appendix 1).

Bedrock and heavy clay soils allow varying degrees of deep root penetration through highly weathered material or through a network of cracks, fissures and channels. Channels, or low resistance pathways, are perma-

nent features of the soil profile, and it has been suggested that they result from dissolution of laterite by humic acid produced by the root itself (Plum and Gosting 1973). Gaiser (1952) found more than 10,000 cavities and root channels per hectare in a hardwood forest in Ohio, United States, pathways that can be reused and expanded by each new generation of trees. Hence, the soil volume should be viewed as a complex network of fissures, cracks and channels on which new root growth largely depends. It has even been suggested that soil compaction in forests may not affect the overall forest productivity, provided that sufficient low resistance pathways allow adequate root development (Nambiar and Sands 1992).

Roots have also been observed penetrating through hard pans and caliche layers in a variety of systems (Silva et al. 1989; Dawson 1993; Day 1994), and into rocks through fissures and cracks (Hellmers et al. 1955, Davis and Pase 1977). Pre-existing old tree channels and earthworm tunnels have also been shown to be important in the downward root development in crop systems (Nambiar and Sands 1992; Nicoulaud et al. 1994).

Finally, some plants find their way deep into the soil by penetrating directly through the bedrock. This phenomenon has been reported for several Mediterranean woody species growing on porous calcareous soils in Israel (Oppenheimer 1958; Orshansky 1951).

### Ecological significance of deep roots

Although a small fraction of root biomass might be found at depths below 1 m, the functional significance of those roots may nevertheless be most important for ecosystem water and carbon fluxes, and nutrient cycling.

The water extracted by plants during the wet season comes from shallow layers where the root density is highest. However, as those layers dry there is a progressive shift towards using deeper water, which allows plants to keep stomata open and extend growth far into the dry season (for review see Gardner 1983). Although we know of the differential water sources in the soil profile, there are very few studies which have quantified the contribution of deep water to the whole ecosystem fluxes. Gregory et al. (1978) showed for winter wheat that few roots below 1 m (about 3% of the total root weight) were responsible for supplying 20% of the transpired water during dry periods. In an Amazonian tropical forest, Nepstad et al. (1994) found that had not considered roots deeper than 2 meters they would have underestimated evapotranspiration by >60% during the dry season. The water available to plants stored below 2 m in the soil provided >75% of the water extracted from the entire soil profile.

There is also plenty of evidence that plants with different rooting habits show different seasonal courses of water potential, and that the duration of water stress and the distribution of soil moisture with depth will deter-

mine whether a species can succeed in a particular environment (Davis and Mooney 1986; Crombie et al. 1988; Sala et al. 1989; Hodgkinson 1992).

For some species (e.g., phreatophytes), survival in arid systems depends exclusively on the capacity to send roots to permanent water tables, as in the case of *Prosopis tamarugo* in the virtually rainless Atacama Desert in Chile (Mooney et al. 1980). Stone and Kalisz (1991) gathered thirty references of plants having contact with water tables at depths from 1.5 to 35 m. In these cases, even if a very small fraction of the roots are tapping water from the water table, the amount of water transferred into the plant may be large. Reicosky et al. (1964) showed that roots tapping water from the water table are hundreds of times more efficient in absorbing it than roots in drier soil. Furthermore, tap roots often show cross sections with a high number of vessels per unit area, indicating a major water transport function (Higgins et al. 1987; see also Pate et al. 1995).

The functional significance of deep roots for water flux in ecosystems under high evaporative conditions has been shown regarding the "hydraulic lift" mechanism which has been reported for several species (Richards and Caldwell 1987; Caldwell and Richards 1989; Dawson 1993). During the night roots take up water from deep soil layers which is released from shallow roots back to the soil in the upper layers. The water is reabsorbed during the next day by the same plants and by shallow-rooted neighbors with no access to deep water. This mechanism has important ecological significance, allowing plants to maintain high transpiration rates during dry periods. Caldwell and Richards (1989) showed that hydraulic lift was responsible for a 30–50% increase of the daytime canopy water flux in artificial mixtures of *Artemisia tridentata* and *Agropyron desertorum* (see also Dawson 1996).

Unlike water relations, much less is known about the contribution of deep soil nutrients to the overall plant nutritional demands. Richter and Markewitz (1995) showed the importance of biological processes in weathering materials in a 8 m-soil profile of a *Pinus taeda* forest in South Carolina, United States; the biological processes were tightly associated with soil influenced by root activity (rhizosphere) all along the soil profile. The importance of deep roots for ecosystem nutrient cycling has also been shown for tropical soils with seasonal drought, Cerrado (Schachtschabel et al. 1992). Nitrate salts from mineralization of organic matter cannot be fully utilized by the vegetation early in the rainy season, and so, are washed out of the top soil down to deep soil horizons. There, nitrate is immobilized by the positive charge balance of  $Fe^{3+}$  and  $Al^{3+}$  found at depths of 1.6 m or more; deep roots will then have access to this nitrate store later in the growing season.

In the deep rhizosphere of *Prosopis glandulosa* in the Chihuahuan desert, United States, a variety of microarthropod taxa has been found down to a depth of 13 m (Silva et al. 1989). The abundance of microarthropods

was positively correlated with root biomass, which suggests that deep rhizosphere processes such as decomposition and mineralization operate in a similar way to those processes in shallow layers. It is also known that plant-feeding nematodes, which are found deep in the rhizosphere, increase nodulation and nitrogen fixation (Huang 1987), and provide infection sites for vesicular-arbuscular mycorrhizal fungi (Freckman and Virginia 1989). In fact, Jenkins et al. (1988) found  $N_2$ -fixing root nodules at a depth of 7 m in the Chihuahuan desert.

Deep roots, in addition to extract water and contribute to the cycling of nutrients, also provide carbon to the soil. In an Amazonian tropical forest Nepstad et al. (1994) found that deep soil layers below 1 m contain large active carbon stocks, 15% of which turns over on annual to decadal timescale. The possession of an active carbon cycle at depth seems to be fairly common in the highly weathered soils in terra firme tropical forest of Amazonia (Trumbore et al. 1995), but almost nothing is known about how common it might be in other biome types.

Ecosystem models which predict carbon sequestration have conventionally used root functional depths between 0.3 m and 2.0 m, which are usually used as fixed factors that do not change or only change for different ecosystem types. The depth at which roots will decay and decompose is essential for determining the ultimate fate of that carbon, and therefore, the capacity of carbon sequestration by different ecosystems. Fisher et al. (1994) showed that increased abundance of introduced deep-rooted grasses in the tropical South American savannas account for an increased sequestration of 100–507 Mt carbon per year, which could explain a substantial part of the missing carbon-sink (Siegenthaler and Sarmiento 1993).

In this review we have shown that deep root habits are quite common in woody and herbaceous species across most of the terrestrial biomes. Roots commonly reach far deeper into the soil than the traditional view has held up to now. This structural trait has important implications for ecosystem water fluxes, as well as for carbon and nutrient cycling, and hence should be appropriately taken into account in the development of ecosystem models.

**Acknowledgements** We acknowledge the generous support by the Max-Planck Forschungspreis through the Alexander von Humboldt Foundation from Germany, NASA-EOS (NAS 5–31726) and NIGEC/DOE (TUL-038–95/96) from the United States. We thank Sabrina Sonntag for providing helpful comments on the manuscript, and Oki Noriko for helping us with the Japanese references. This work contributes to the Global Change and Terrestrial Ecosystems (GCTE) Core Project of the International Geosphere-Biosphere Program (IGBP).

**Appendix 1** Reported maximum rooting depth (m) by species with soil type, country and reference grouped by biome

Species/ dominant species <sup>a</sup>	Maximum rooting depth (m)	Soil type	Country	Reference
<b>BOREAL FOREST</b>				
<i>Larix laricina</i>	1.2	medium-coarse sand/podzol	S-Canada	Bannan 1940
<i>Larix sibirica</i>	1.8	medium-loamy	Russia	Verzunov 1980
<i>Picea glauca</i>	1.8	medium-loamy	Russia	Verzunov 1980
<i>Pinus banksia</i>	1.2	medium-coarse sand/podzol	S-Canada	Bannan 1940
<i>Pinus banksiana</i>	2.0	aeolian sands/Eutric brunisol	S-Canada	Strong and La Roi 1983
<i>Pinus contorta</i>	3.3	–	S-Canada	Horton 1958
<i>Populus tremuloides</i>	2.0	sandy substrate	S-Canada	Strong and La Roi 1983
<b>CROPS</b>				
<i>Andropogon sorghum</i>	1.1	lowland silt loam	Nebraska, USA	Weaver 1926
<i>Avena sativa</i>	1.8	–	Kansas, USA	Weaver 1926
<i>Beta vulgaris</i>	1.8	sandy loam	Nebraska, USA	Weaver 1926
<i>Bromus inermis</i>	1.1	silty-clay to clay-loam alluvial	Canada	Leyshon 1991
<i>Elymus angustus</i>	3.5	–	S-Canada	Lawrence 1975
<i>Elymus junceus</i>	1.8	–	S-Canada	Lawrence 1975
<i>Glycine max</i>	1.8	Muir silt loam	Kansas, USA	Mayaki et al. 1976
<i>Helianthus annuus</i>	2.7	lowland silt loam	Nebraska, USA	Weaver 1926
<i>Helianthus annuus</i>	2.7	Muir silt loam from alluvium	Kansas, USA	Jaafar et al. 1993
<i>Hordeum vulgare</i>	1.3	Lowland silt loam	Nebraska, USA	Weaver 1926
<i>Hordeum sp.</i>	2.2	loamy sand/Xeric Psamment	W-Australia	Hamblin and Tennant 1987
<i>Lupinus angustifolius</i>	2.5	loamy sand/Xeric Psamment	W-Australia	Hamblin and Tennant 1987
<i>Medicago sativa</i>	3.7	–	Nebraska, USA	Weaver 1926
<i>Secale cereale</i>	1.5	silt loam/hard clayey subsoil	Nebraska, USA	Weaver 1926
<i>Solanum tuberosum</i>	1.4	mellow loess soil	Nebraska, USA	Weaver 1926
<i>Triticum aestivum</i>	1.0	–	England	Welbank et al. 1974
<i>Triticum aestivum</i>	1.4	Muir silt loam	Kansas, USA	Chaudhuri et al. 1990
<i>Triticum aestivum</i>	1.5	lowland silt loam	Nebraska, USA	Weaver 1926
<i>Triticum aestivum</i>	1.8	–	Mid Europe	Kutschera 1960
<i>Triticum aestivum</i>	3.0	loamy sand/Xeric Psamment	W-Australia	Hamblin and Tennant 1987
<i>Triticum durum</i>	2.3	loess soil	Nebraska, USA	Weaver 1926
<i>Zea mays</i>	1.3	deep clay loam	E-France	Pages and Pellerin 1994
<i>Zea mays</i>	2.4	–	Nebraska, USA	Weaver 1926
<b>DESERT</b>				
<i>Alhagi maurorum</i>	15.0	river banks	Israel	Shmueli 1948
<i>Artemisia monosperma</i>	5.0	sand dunes	Israel	Zohary and Fahn 1952
<i>Artemisia tridentata</i>	1.8	shale/sandstone bedrock	Colorado, USA	Branson et al. 1976
<i>Artemisia tridentata</i>	2.2	loamy-skeletal/Haploxerolls	Utah, USA	Richards and Caldwell 1987
<i>Artemisia tridentata</i>	2.3	aeolian sandy loam	Idaho, USA	Reynolds and Fraley 1989
<i>Atriplex halimus</i>	8.0	alluvia soils/run-on habitats	Israel	Zohary 1961
<i>Chrysothamnus vicidiflorus</i>	2.0	aeolian sandy loam	Idaho, USA	Reynolds and Fraley 1989
<i>Franseria deltoidea</i>	1.8	wash with hardpan (caliche)	Arizona, USA	Cannon 1911
<i>Hammada salicornica</i>	2.5	sand dunes	Israel	Zohary and Orshan 1949
<i>Leptadenia pyrotechnica</i>	11.5	sandy and silty/clay at depth	Egypt	Batanouny and Wahab 1973
<i>Leymus cinereus</i>	2.0	aeolian sandy loam	Idaho, USA	Reynolds and Fraley 1989
<i>Mulinum spinosum<sup>a</sup></i>	3.0	fine sand/caliche layer at 0.6 m	S-Argentina	Schulze et al. 1996
<i>Nassauvia glomerulosa<sup>a</sup></i>	3.0	sandy loam/caliche at 0.7 m	S-Argentina	Schulze et al. 1996
<i>Nitraria retusa</i>	5.0	sandy	Israel	Ginzburg 1966
<i>Ochradenus baccatus</i>	5.0	sandy	Israel	Ginzburg 1966
<i>Prosopis farcta</i>	15.0	river banks	Israel	Zohary and Orshan 1949
<i>Prosopis glandulosa</i>	2.0	Nuvalde clay loam	Texas, USA	Heitschmidt et al. 1988
<i>Prosopis glandulosa</i>	6.0	clay loam/sand, clay at depth	California, USA	Nilsen et al. 1983
<i>Prosopis glandulosa</i>	12.0	sandy/Torrifluent	New Mexico, USA	Freckman and Virginia 1989
<i>Prosopis glandulosa</i>	15.0	clay loam	New Mexico, USA	Silva et al. 1989
<i>Prosopis juliflora</i>	53.0	–	Arizona, USA	Phillips 1963
<i>Prosopis tamarugo</i>	3.5	–	N-Chile	Mooney et al. 1980
<i>Prosopis velutina</i>	8.0	–	Arizona, USA	Cannon 1911
<i>Retama raetam</i>	20.0	sand dunes	Israel	Zohary and Fahn 1952
<i>Tamarix aphylla</i>	20.0	alluvial soils/run-on habitats	Israel	Zohary 1961
<i>Tamarix pentantra</i>	3.6	alluvial banks	Arizona, USA	Gary 1963
<i>Zilla spinosa</i>	5.0	–	Israel	Ginzburg 1966

## Appendix 1 (continued)

Species/ dominant species <sup>a</sup>	Maximum rooting depth (m)	Soil type	Country	Reference
<b>SCLEROPHYLLOUS SHRUBLAND AND FOREST</b>				
<i>Shrubs</i>				
<i>Adenostoma fasciculatum</i>	2.4	silt sandy	California, USA	Hanes 1965
<i>Adenostoma fasciculatum</i>	7.6	sandy loam on anorthosite	California, USA	Hellmers et al. 1955
<i>Adenostoma sparsifolium</i>	2.4	silt sandy	California, USA	Hanes 1965
<i>Arbutus unedo</i>	3.5	sandy loam	NE Spain	J Canadell, unpublished work
<i>Arctostaphylos glandulosa</i>	5.2	sandy loam on granodiorite	California, USA	Hellmers et al. 1955
<i>Arctostaphylos glauca</i>	2.6	sandy loam on granodiorite	California, USA	Hellmers et al. 1955
<i>Arctostaphylos glutinosa</i>	2.5	shallow on fractured shales	California, USA	Davis 1972
<i>Arctostaphylos pallida</i>	4.0	shallow on fractured shales	California, USA	Davis 1972
<i>Banksia marginata</i>	2.4	sandy	SE-Australia	Specht and Rayson 1957
<i>Banksia ornata</i>	2.4	sandy	SE-Australia	Specht and Rayson 1957
<i>Baccharis pilularis</i>	3.2	packed sand like a rock	California, USA	Wright 1928
<i>Banksia spp.</i>	5.0	podsolized sand	SW-Australia	Low and Lamont 1990
<i>Calytrix flavescens</i>	2.0	grey sands with hardpan	SW-Australia	Dodd et al. 1984
<i>Casuarina muelleriana</i>	2.0	sandy	SE-Australia	Specht and Rayson 1957
<i>Casuarina pusilla</i>	2.4	sandy	SE-Australia	Specht and Rayson 1957
<i>Ceanothus leucodermis</i>	3.7	sandy loam on granodiorite	California, USA	Hellmers et al. 1955
<i>Ceanothus megacarpus</i>	2.4	sandstone with fissures	California, USA	Thomas and Davis 1989
<i>Ceanothus oliganthus</i>	1.8	clay loam on diorite	California, USA	Hellmers et al. 1955
<i>Ceanothus spinosus</i>	3.1	sandstone with fissures	California, USA	Thomas and Davis 1989
<i>Daviesia brevifolia</i>	2.0	sandy	SE-Australia	Specht and Rayson 1957
<i>Eremaea beaufortoides</i>	6.0	alluvial sand with colluvium	SW-Australia	Hnatiuk and Hopkins 1980
<i>Eremaea pauciflora</i>	2.4	grey sands with hardpan	SE-Australia	Dodd et al. 1984
<i>Erica arborea</i>	2.0	sandy loam	NE-Spain	J. Canadell, unpublished work
<i>Hibbertia hypericoides</i>	2.1	grey sands with hardpan	SW-Australia	Dodd et al. 1984
<i>Jacksonia floribunda</i>	3.1	grey sands with hardpan	SW Australia	Dodd et al. 1984
<i>Jacksonia furcellata</i>	2.0	grey sands with hardpan	SW Australia	Dodd et al. 1984
<i>Laudonia behrii</i>	2.0	sandy	SE-Australia	Specht and Rayson 1957
<i>Leptospermum myrsinoides</i>	2.3	sandy	SE-Australia	Specht and Rayson 1957
<i>Leucadendron salignum</i>	3.0	loamy medium sand	South Africa	Higgins et al. 1987
<i>Lithraea caustica</i>	5.0	-	Central Chile	Gilberto and Estay 1978
<i>Melaleuca scabra</i>	2.0	grey sands with hardpan	SW-Australia	Dodd et al. 1984
<i>Melaleuca seriata</i>	2.1	grey sands with hardpan	SW-Australia	Dodd et al. 1984
<i>Petrophile linearis</i>	2.0	grey sands with hardpan	SW-Australia	Dodd et al. 1984
<i>Photinia arbutifolia</i>	2.1	clay loam on diorite	California, USA	Hellmers et al. 1955
<i>Phyllota pleurandroides</i>	2.3	sandy	SE-Australia	Specht and Rayson 1957
<i>Phyllota remota</i>	2.4	sandy	SE-Australia	Specht and Rayson 1957
<i>Protea nerifolia</i>	3.0	loamy medium sand	South Africa	Higgins et al. 1987
<i>Protea repens</i>	3.0	loamy medium sand	South Africa	Higgins et al. 1987
<i>Quercus calliprinos<sup>a</sup></i>	4.6	terra-rossa on limestone	Israel	Shachori et al. 1967
<i>Quercus dumosa</i>	8.5	clay loam on diorite	California, USA	Hellmers et al. 1955
<i>Quercus turbinella</i>	6.4	fracturated granite	Arizona, USA	Davis and Pase 1977
<i>Quercus turbinella</i>	9.1	alluvial and redish brown	Arizona, USA	Saunier and Wagle 1967
<i>Quillaja saponaria</i>	8.0	-	Central Chile	Gilberto and Estay 1978
<i>Rhus glabra</i>	6.7	loess hills	Nebraska, USA	Weaver 1919
<i>Rhus laurina</i>	5.4	sandstone with fissures	California, USA	Thomas and Davis 1989
<i>Rhus laurina</i>	13.2	-	California, USA	DeSouza et al. 1986
<i>Salvia apiana</i>	1.5	coarse, loose gravel	California, USA	Hellmers et al. 1955
<i>Scholtzia involucreta</i>	1.9	grey sands with hardpan	SW-Australia	Dodd et al. 1984
<i>Spyridium subochreatum</i>	1.9	sandy	SE-Australia	Specht and Rayson 1957
<i>Stirlingia latifolia</i>	2.6	grey sands with hardpan	SW-Australia	Dodd et al. 1984
<i>Xanthorrhoea australis</i>	2.4	sandy	SE-Australia	Specht and Rayson 1957
<i>Trees</i>				
<i>Eucalyptus marginata</i>	15.0	lateritic, sandy-clay at depth	SW-Australia	Kimber 1974
<i>Eucalyptus marginata</i>	20.0	-	SW-Australia	Carbon et al. 1980
<i>Eucalyptus marginata</i>	40.0	fissured granite, clay subsoil	SW-Australia	Dell et al. 1983
<i>Eucalyptus regnans</i>	2.7	-	SW-Australia	Incoll 1969
<i>Eucalyptus signata</i>	3.0	sandy	NE-Australia	Westman and Rogers 1977
<i>Eucalyptus sp.</i>	10.0	sand dunes	NE-Australia	Westman and Rogers 1977
<i>Quercus agrifolia</i>	10.7	-	California, USA	Cannon 1914
<i>Quercus chrysolepis</i>	7.3	sandy loam on granodiorite	California, USA	Hellmers et al. 1955
<i>Quercus douglasii</i>	3.7	alluvial loam	California, USA	Cannon 1914
<i>Quercus ilex</i>	3.7	sandstone	NE-Spain	J. Canadell, unpublished work
<i>Quercus wislizenii<sup>a</sup></i>	22.9	fractured rock	California, USA	Lewis and Burgy 1964

## Appendix 1 (continued)

Species/ dominant species <sup>a</sup>	Maximum rooting depth (m)	Soil type	Country	Reference
<b>TEMPERATE CONIFEROUS FOREST</b>				
<i>Abies firma</i>	3.3	sandy soil	Japan	Karizumi 1979
<i>Picea excelsa</i>	2.3	silt loam	Japan	Karizumi 1979
<i>Pinus densiflora</i>	3.4	silt loam	Japan	Karizumi 1979
<i>Pinus echinata</i>	3.3	sandy soil	New Jersey, USA	Lull and Axley 1958
<i>Pinus elliotii</i>	3.3	—	Florida, USA	van Rees and Comerford 1986
<i>Pinus halepensis</i> <sup>a</sup>	7.3	terra-rossa on limestone	Israel	Shachori et al. 1967
<i>Pinus halepensis</i>	7.5	weathered granite	NE-Spain	J. Canadell, unpublished work
<i>Pinus luchuensis</i>	3.5	sandy loam	Japan	Karizumi 1979
<i>Pinus palustris</i>	4.8	Norfold sand deep phase	Florida, USA	Heyward 1933
<i>Pinus pinaster</i>	7.0	—	Australia	Butcher and Havel 1976
<i>Pinus pinea</i>	5.0	weathered granite	NE-Spain	J. Canadell, unpublished work
<i>Pinus ponderosa</i>	3.5	clay loam soil	Oregon, USA	Zwieniecki and Newton 1994
<i>Pinus radiata</i>	2.0	sandy soil	S-Australia	Nambiar and Sands 1992
<i>Pinus resinosa</i>	2.7	Hinckley coarse sand	New York, USA	White and Wood 1958
<i>Pinus resinosa</i>	5.0	sandy outwash	New York, USA	Leaf et al. 1955
<i>Pinus rigida</i>	2.7	sandy soil	New Jersey, USA	McQuilkin 1935
<i>Pinus rigida</i>	3.4	sandy soil	Japan	Karizumi 1979
<i>Pinus strobus</i>	2.8	sandy soil	Japan	Karizumi 1979
<i>Pinus sylvestris</i>	2.7	sand overlying chalky drift	United Kingdom	Roberts 1976
<i>Pinus taeda</i>	2.0	fullerton and bodine	Tennessee, USA	Harris et al. 1977
<i>Pinus taeda</i>	4.0	granite wheathered/Ultisol	S-Carolina, USA	Richter and Markewitz 1995
<b>TEMPERATE DECIDUOUS FOREST</b>				
<i>Acer negundo</i>	4.0	upland clay	Missouri, USA	Biswell 1935
<i>Acer saccharum</i>	3.7	silty loams with hardpan	New York, USA	Dawson 1993
<i>Carya spp.</i>	1.8	sandstone	Ohio, USA	Gaiser 1952
<i>Corylus americana</i>	3.5	loess hills	Nebraska, USA	Weaver 1919
<i>Fraxinus japonica</i>	2.0	fine texture clay	Japan	Karizumi 1979
<i>Juglans nigra</i>	3.0	silt loam	Japan	Karizumi 1979
<i>Latrix decidua</i>	3.4	fine silty sand at depth	New York, USA	White and Wood 1958
<i>Nothofagus pumila</i>	2.0	orange loam/rocks at depth	S-Argentina	Schulze et al. 1996
<i>Platanus orientalis</i>	2.6	medium texture	Japan	Karizumi 1979
<i>Populus nigra</i>	1.9	silt loam	Japan	Karizumi 1979
<i>Populus sargentii</i>	2.6	loam underlain with clay	Missouri, USA	Biswell 1935
<i>Populus tremula</i>	2.0	clay subsoil	Sweden	Persson 1975
<i>Populus tremuloides</i>	2.3	grey clay	Michigan, USA	Day 1944
<i>Populus tremuloides</i>	2.9	sandy loam	Utah, USA	Gifford 1966
<i>Prunus yedoensis</i>	2.1	fine texture clay	Japan	Karizumi 1979
<i>Quercus dentata</i>	4.3	silt loam	Japan	Karizumi 1979
<i>Quercus macrocarpa</i>	4.3	fine-textured loams	Nebraska, USA	Weaver and Kramer 1932
<i>Quercus macrocarpa</i>	4.4	upland clay	Missouri, USA	Biswell 1935
<i>Quercus sp.-Carya sp.</i> <sup>a</sup>	4.0	silt loam on sandstone/shale	Virginia, USA	Kochenderfer 1973
<i>Quercus velutina</i>	3.0	medium texture	Japan	Karizumi 1979
<i>Salix babylonica</i>	2.2	silt loam	Japan	Karizumi 1979
<b>TEMPERATE GRASSLAND</b>				
<i>Agropyron repens</i>	2.4	loose sandy	Nebraska, USA	Weaver 1919
<i>Agropyron smithii</i>	2.7	silt loam	Colorado, USA	Weaver 1958
<i>Agropyron spicatum</i>	1.4	med. textur. Benge series	Washington, USA	Harris 1967
<i>Agropyron spicatum</i>	1.5	silt loam	Washington, USA	Weaver 1919
<i>Amorpha canescens</i>	5.0	loose sandy	Nebraska, USA	Weaver 1919
<i>Andropogon furcatus</i>	1.5	Judson silt loam	Nebraska, USA	Weaver and Darland 1949
<i>Andropogon furcatus</i>	2.8	clay loam	Nebraska, USA	Weaver 1919
<i>Andropogon gerardi</i>	2.1	lilt loam	Iowa, USA	Weaver 1958
<i>Andropogon hallii</i>	1.8	sandy	Nebraska, USA	Tolstead 1942
<i>Andropogon hallii</i>	3.0	sandy	Colorado, USA	Weaver 1958
<i>Andropogon scoparius</i>	1.5	silt loam	Iowa, USA	Weaver 1958
<i>Andropogon scoparius</i>	1.8	loam sandy	Colorado, USA	Weaver 1919
<i>Aragallus lambertii</i>	1.4	loam sandy	Colorado, USA	Weaver 1919
<i>Argemone platyceras</i>	3.7	loam sandy	Colorado, USA	Weaver 1919
<i>Artemisia frigida</i>	1.7	dark brown soil on shales	S-Canada	Coupland and Johnson 1965
<i>Artemisia cana</i>	2.4	dark brown soil on shales	S-Canada	Coupland and Johnson 1965
<i>Atriplex nuttallii</i>	1.8	dark brown soil on shales	S-Canada	Coupland and Johnson 1965
<i>Astragalus crassicaarpus</i>	2.0	loam soil on hard joint clay	Nebraska, USA	Weaver 1919
<i>Berberis repens</i>	3.0	silt loam	Washington, USA	Weaver 1919
<i>Biscutella laevigata</i>	2.1	—	Mid Europe	Kutschera 1960



## Appendix 1 (continued)

Species/ dominant species <sup>a</sup>	Maximum rooting depth (m)	Soil type	Country	Reference
<i>Bouteloua curtipendula</i>	1.7	silt loam	Colorado, USA	Weaver 1958
<i>Bouteloua gracilis</i>	1.7	Colby silt loam	Nebraska, USA	Weaver and Darland 1949
<i>Bouteloua gracilis</i>	1.8	silt loam	Colorado, USA	Weaver 1958
<i>Bouteloua gracilis</i> <sup>a</sup>	2.1	–	Kansas, USA	Albertson et al. 1953
<i>Brauneria pallida</i>	2.4	clay loam	Nebraska, USA	Weaver 1919
<i>Buchloe dactyloides</i>	1.8	silt loam	Iowa, USA	Weaver 1958
<i>Buchloe dactyloides</i>	2.0	Wabash silt loam	Nebraska, USA	Weaver and Darland 1949
<i>Bulbilis dactyloides</i>	1.9	alluvial	Nebraska, USA	Weaver 1919
<i>Calamovilfa longifolia</i>	1.8	dark brown soil on shales	S-Canada	Coupland and Johnson 1965
<i>Calamovilfa longifolia</i>	3.0	sandy	Colorado, USA	Weaver 1958
<i>Carex arenaria</i>	1.8	–	Mid Europe	Kutschera 1960
<i>Carex filifolia</i>	1.5	silt loam	Colorado, USA	Weaver 1958
<i>Carlina acaulis</i>	4.1	–	Germany	Kutschera 1960
<i>Centaurea scabiosa</i>	3.3	–	Germany	Kutschera 1960
<i>Chrysopsis villosa</i>	2.4	dark brown soil on shales	S-Canada	Coupland and Johnson 1965
<i>Equisetum arvense</i>	3.0	sandy	Canada	Coupland and Johnson 1965
<i>Equisetum palustre</i>	2.5	–	Mid Europe	Kutschera 1960
<i>Eriogonum heracleoides</i>	2.4	silt loam	Washington, USA	Weaver 1919
<i>Eriogonum jamesii</i>	2.3	loam with some sand	Colorado, USA	Weaver 1919
<i>Eriogonum microthecum</i>	3.0	sandy	Colorado, USA	Weaver 1919
<i>Erodium botrys</i>	1.3	gravelly clay loam	California, USA	McKell et al. 1962
<i>Eryngium campestre</i>	4.2	–	Germany	Kutschera 1960
<i>Eurotia lanata</i>	1.8	dark brown soil on shales	S-Canada	Coupland and Johnson 1965
<i>Festuca arizonica</i>	1.2	sandy loam-sandy clay	Colorado, USA	Currie and Hammer 1979
<i>Festuca arizonica</i>	1.3	–	Colorado, USA	Schuster 1964
<i>Festuca arundinacea</i>	2.7	–	Germany	Kutschera 1960
<i>Festuca pallescens</i> <sup>a</sup>	2.0	alluvial sandy loam & gravel	S-Argentina	Schulze et al. 1996
<i>Gaillardia aristata</i>	1.7	dark brown soil on shales	S-Canada	Coupland and Johnson 1965
<i>Geranium viscosissimum</i>	2.9	silt loam	Washington, USA	Weaver 1919
<i>Grindelia squarrosa</i>	1.9	loose sand	Nebraska, USA	Weaver 1919
<i>Heracleum sphondyleum</i>	2.0	–	Mid Europe	Kutschera 1960
<i>Hieracium scouleri</i>	2.2	silt loam	Washington, USA	Weaver 1919
<i>Hoorebekia racemosa</i>	3.4	silt loam	Washington, USA	Weaver 1919
<i>Kochia prostrata</i>	6.3	–	Germany	Kutschera 1960
<i>Kuhnia glutinosa</i>	5.2	–	Nebraska, USA	Weaver 1919
<i>Lepachys pinnata</i>	1.5	brown silt loam	Illinois, USA	Sperry 1935
<i>Lespedeza capitata</i>	2.4	lower slopes of loess hills	Nebraska, USA	Weaver 1919
<i>Liatris punctata</i>	2.1	gravelly	S-Canada	Coupland and Johnson 1965
<i>Liatris punctata</i>	4.8	clay	Nebraska, USA	Weaver 1919
<i>Lithospermum gmelini</i>	2.1	sandy	Nebraska, USA	Tolstead 1942
<i>Lupinus ornatus</i>	4.0	silt loam	Washington, USA	Weaver 1919
<i>Lygodesmia juncea</i>	3.0	sandy	Nebraska, USA	Tolstead 1942
<i>Lygodesmia juncea</i>	3.0	dark brown soil on shales	S-Canada	Coupland and Johnson 1965
<i>Lygodesmia juncea</i>	6.3	loess	Nebraska, USA	Weaver 1919
<i>Medicago falcata</i>	4.3	–	Germany	Kutschera 1960
<i>Muhlenbergia montana</i>	1.3	sandy clay loam subsoil	Colorado, USA	Schuster 1964
<i>Onobrychis natrix</i>	2.3	–	Mid Europe	Kutschera 1960
<i>Ononis natrix</i>	2.3	–	Germany	Kutschera 1960
<i>Panicum virgatum</i>	2.7	loose sand	Nebraska, USA	Weaver 1919
<i>Parthenium integrifolium</i>	1.8	brown silt loam	Illinois, USA	Sperry 1935
<i>Petalostemum purpureum</i>	1.8	brown silt loam	Illinois, USA	Sperry 1935
<i>Peucedanum cervaria</i>	4.1	–	Germany	Kutschera 1960
<i>Phalaris aquatica</i>	1.2	granite	Spain	Joffre et al. 1987
<i>Pimpinella saxifraga</i>	3.7	–	Germany	Kutschera 1960
<i>Potentilla blaschkeana</i>	2.3	silt loam	Washington, USA	Weaver 1919
<i>Potentilla fruticosa</i>	3.0	gravelly	S-Canada	Coupland and Johnson 1965
<i>Potentilla concinna</i>	1.8	dark brown soil on shales	S-Canada	Coupland and Johnson 1965
<i>Psoralea tenuiflora</i>	1.8	loose sand	Nebraska, USA	Weaver 1919
<i>Psoralea tenuiflora</i>	3.7	loam sandy, silt loam	Colorado, USA	Weaver 1919
<i>Redfieldia flexuosa</i>	1.5	sandy	Colorado, USA	Weaver 1958
<i>Ruellia ciliosa</i>	1.5	brown silt loam	Illinois, USA	Sperry 1935
<i>Rumex crispus</i>	3.3	–	Germany	Kutschera 1960
<i>Senecio riddellii</i>	1.5	sandy soil	Nebraska, USA	Tolstead 1942
<i>Silphium integrifolium</i>	1.7	brown silt loam	Illinois, USA	Sperry 1935
<i>Silphium laciniatum</i>	1.8	brown silt loam	Illinois, USA	Sperry 1935
<i>Solidago canadensis</i>	3.4	loose sandy	Nebraska, USA	Weaver 1919
<i>Solidago rigida</i>	1.4	brown silt loam	Illinois, USA	Sperry 1935
<i>Spartina pectinata</i>	4.0	silt loam	Iowa, USA	Weaver 1958

## Appendix 1 (continued)

Species/ dominant species <sup>a</sup>	Maximum rooting depth (m)	Soil type	Country	Reference
<i>Sporobolus cryptandrus</i>	1.5	deeply eroded loess	Nebraska, USA	Weaver 1919
<i>Sporobolus heterolepis</i>	1.5	silt loam	Iowa, USA	Weaver 1958
<i>Stipa spartea</i>	1.8	silt loam	Iowa, USA	Weaver 1958
<i>Taraxacum serotinum</i>	4.6	–	Germany	Kutschera 1960
<i>Thermopsis rhombifolia</i>	2.1	gravelly	S-Canada	Coupland and Johnson 1965
<i>Tradescantia reflexa</i>	1.6	brown	Illinois, USA	Sperry 1935
<i>Vernonia baldwinii</i>	3.5	loose sand	Nebraska, USA	Weaver 1919
<b>TROPICAL DECIDUOUS FOREST</b>				
<i>Antiaris toxicaria</i>	3.5	red soil	China	Bang-Xing 1991
<i>Baccaurea ramiflora</i>	3.7	red soil	China	Bang-Xing 1991
<i>Gironniera subaequalis</i>	4.7	red soil	China	Bang-Xing 1991
<i>Symplocos cochinchinensis</i>	2.0	red soil	China	Bang-Xing 1991
<i>Xanthophyllum siamense</i>	4.6	red soil	China	Bang-Xing 1991
<b>TROPICAL EVERGREEN FOREST</b>				
<i>Apodytes dimidiata</i>	8.2	sandy loam on schists	Kenya	Kerfoot 1963
<i>Chlorophora excelsa</i>	2.0	ferralitic	Ghana	Mensah and Jenik 1968
<i>Chlorophora excelsa</i>	3.0	ferralitic	Ghana	Jenik 1971
Community	18.0	clay	Brazil	Nepstad et al. 1994
Community	5.0	–	Brazil	Poels 1987
Community	2.5	Turraeantho on sandy soil	Ivory Coast	Huttel 1975
<b>TROPICAL GRASSLAND AND SAVANNA</b>				
<i>Acacia erioloba</i>	60.0	Kalahari sands	Botswana	Jennings 1974
<i>Anacardium pumilum</i>	10.0	–	Brazil	Ferri 1961
<i>Andira humilis</i>	18.0	reddish loamy earth	Brazil	Rawitscher 1948
<i>Andira spp.</i>	19.0	–	Brazil	Rawitscher et al. 1943
<i>Aristolachia giberti</i>	1.8	reddish loamy earth	Brazil	Rawitscher 1948
<i>Boscia albitrunca</i>	68.0	Kalahari sands	Botswana	Jennings 1974
<i>Brachiaria brizantha</i> <sup>a</sup>	8.0	clay	Brazil	Nepstad et al. 1994
<i>Brachystegia sp.</i>	1.8	–	Zimbabwe	Strang 1969
<i>Capparis sp.</i>	1.6	sandy clay loam	Ghana	Okali et al. 1973
<i>Curatella americana</i>	4.0	–	Venezuela	Foldats and Rutkis 1975
<i>Jacaranda decurrens</i> <sup>a</sup>	11.0	–	Brazil	Rawitscher et al. 1943
<i>Ochna pulchra</i>	2.2	structureless sand	South Africa	Rutherford 1983
<i>Panicum maximum</i> <sup>a</sup>	12.0	clay	Brazil	Nepstad et al. 1994
<i>Stipagrostis amabilis</i> <sup>a</sup>	5.0	Kalahari sands	South Africa	J. Canadell, unpublished work
<i>Stryphnodendron sp.</i>	2.0	reddish loamy earth	Brazil	Rawitscher 1948
<b>TUNDRA</b>				
<i>Cares aquatilis</i> <sup>a</sup>	0.3	–	Alaska, USA	Dennis et al. 1978
<i>Dryas punctata</i> <sup>a</sup>	0.5	permafrost at 40–55 cm	N-Russia	Khodachek 1971
<i>Dupontia fischeri</i> <sup>a</sup>	0.3	organic matter on sediments	Alaska, USA	Dennis 1977
<i>Eriophorum vaginatum</i> <sup>a</sup>	0.6	silty soil on permafrost	Alaska, USA	Wein and Bliss 1974
<i>Betula nana</i>	0.5	permafrost at 50 cm	Alaska, USA	S. Hobbie, unpublished work
<i>Luzula confusa</i>	0.3	loams	N-Canada	Bliss and Svoboda 1984
<i>Salix glauca</i>	0.5	permafrost at 45–60 cm	W-Russia	Ignatenko and Khakimzy 1971
<i>Salix planifolia</i>	0.9	coarse textured/bottom pit	Colorado, USA	Webber and May 1977

<sup>a</sup> Maximum rooting depth is not linked to the species name but to the dominant species in the community

## References

- Albertson FW, Riegel A, Launchbaugh JL (1953) Effects of different intensities of clipping on short grasses in west-central Kansas. *Ecology* 34:1–20
- Bang-Xing W (1991) Studies on the vertical structure of seasonal rain-forest in Xishuangbanna of Yunnan. *Acta Bot Sin* 33: 232–239
- Bannan (1940) The root systems of northern Ontario conifers growing in sand. *Am J Bot* 27: 108–114
- Batanouny KH, Abdel Wahab AM (1973) Eco-physiological studies on desert plants. VIII. Root penetration of *Leptadenia pyrotechnica* (Forsk.) Decne. in relation to its water balance. *Oecologia* 11: 151–161
- Bennie ATP (1991) Growth and mechanical impedance. In: Waisel Y, Eshel A, Kafkafi U (eds) *Plant roots: the hidden half* (Books in soils, plants, and the environment). Marcel Dekker, New York, pp 393–414
- Biswell HH (1935) Effects of environment upon the root habits of certain deciduous forest trees. *Bot Gaz* 9: 676–708
- Bliss LC, Svoboda J (1984) Plant communities and plant production in the western Queen Elizabeth Islands. *Holarct Ecol* 7: 325–344
- Branson FA, Miller RF, McQueen IS (1976) Moisture relationships in twelve northern desert shrub communities near Grand Junction, Colorado. *Ecology* 57: 1104–1124
- Butcher TB, Havel JJ (1976) Influence of moisture relationships on thinning practice. *NZ J For Sci* 6: 158–170

- Caldwell MM, Richards JH (1989) Hydraulic lift: water efflux from upper roots improves effectiveness of water uptake by deep roots. *Oecologia* 79: 1–5
- Canadell J, Zedler PH (1995) Underground structures of woody plants in Mediterranean ecosystems of Australia, California, and Chile. In: Arroyo MTK, Zedler PH, Fox MD (eds) *Ecology and biogeography of mediterranean ecosystems in Chile, California, and Australia*. Springer, Berlin Heidelberg New York, pp 177–210
- Cannon WA (1911) The root habits of desert plants. *Carnegie Inst Wash Publ* 131: 1–96
- Cannon WA (1914) Specialization in vegetation and in environment in California. *Plant World* 17: 223–237
- Carbon BA, Bartle GA, Murray AM, MacPherson DK (1980) The distribution of root length, and the limits to flow of soil water to roots in a dry sclerophyll forest. *For Sci* 26: 656–664
- Chaudhuri UN, Kirkham MB, Kanemasu ET (1990) Root growth of winter wheat under elevated carbon dioxide and drought. *Crop Sci* 30: 853–857
- Coupland RT, Johnson RE (1965) Rooting characteristics of native grassland species in Saskatchewan. *J Ecol* 53: 475–507
- Crombie DS, Tippet JT, Hill TC (1988) Down water potential and root depth of trees and understory species in southwestern Australia. *Aust J Bot* 36: 621–631
- Currie PO, Hammer FL (1979) Detecting depth and lateral spread of roots of native range plants using radioactive phosphorus. *J Range Manage* 32: 101–103
- Davis CB (1972) Comparative ecology of six members of the *Arcostaphylos andersonii* complex. PhD dissertation, University of California, Davis
- Davis SD, Mooney HA (1986) Water use patterns of four co-occurring chaparral shrubs. *Oecologia* 70: 172–177
- Davis EA, Pase CP (1977) Root system of shrub live oak: implications for water yield in Arizona chaparral. *J Soil Water Conserv* 32: 174–180
- Dawson TE (1993) Hydraulic lift and water use by plants: implications for water balance, performance and plant-plant interactions. *Oecologia* 95: 565–574
- Dawson TE (1996) Determining water use by trees and forests from isotopic, energy balance and transpiration analyses: the roles of tree size and hydraulic lift. *Tree Physiol* 16: 263–272
- Day MW (1944) The root system of aspen. *Am Midl Nat* 32: 502–509
- Dell B, Bartle JR, Tacey WH (1983) Root occupation and root channels of jarrah forest subsoils. *Aust J Bot* 31: 615–627
- Dennis JG (1977) Distribution patterns of belowground standing crop in arctic tundra at Barrow, Alaska. In: Marshall JK (ed) *The belowground ecosystem: a synthesis of plant-associated processes*. Range Science Department, Colorado State University, Fort Collins
- Dennis JG, Tieszen LL, Vetter MA (1978) Seasonal dynamics of above- and belowground production of vascular plants at Barrow, Alaska. In: Tieszen LL (ed) *Vegetation and production ecology of the Alaskan Arctic tundra (Ecological studies 29)*. Springer, Berlin Heidelberg New York, pp 113–140
- DeSouza J, Silka PA, Davis SD (1986) Comparative physiology of burned and unburned *Rhus laurina* after chaparral wildfire. *Oecologia* 71: 63–68
- Dodd J, Heddle EM, Pate EM, Dixon KW (1984) Rooting patterns of sandplain plants and their functional significance. In: Pate JS, Beard JS (eds) *Kwongan: plant life of the sandplain*. University of Western Australia Press, Nedlands, pp 146–177
- Feldman LJ (1984) Regulation of root development. *Annu Rev Plant Physiol* 35: 223–242
- Ferri MG (1961) Problems of water relations of some Brazilian vegetation types, with special consideration of the concepts of xeromorphy and xerophytism. Plant-water relationships in arid and semi-arid continents. In: UNESCO (ed) *Arid zone research 16*. UNESCO, Paris, pp 191–197
- Fisher MJ, Rao IM, Ayarza MA, Lascano CE, Sanz JI, Thomas RJ, Vera RR (1994) Carbon storage by introduced deep-rooted grasses in the South American savannas. *Nature* 371: 236–238
- Foldats E, Rutkis E (1975) Ecological studies of chaparral (*Curatella americana* L.) and manteco (*Byrsonima crassifolia* H.B.K.) in Venezuela. *J Biogeogr* 2: 159–178
- Freckman DW, Virginia RA (1989) Plant-feeding nematodes in deep-rooting desert ecosystems. *Ecology* 70: 1665–1678
- Gardner WR (1983) Soil properties and efficient water use: an overview. In: Taylor HM, Jordan WR, Sinclair TR (eds) *Limitations to efficient water use in crop production*. American Society of Agronomy, Madison, pp 45–64
- Gaiser RN (1952) Root channels and roots in forest soils. *Soil Sci Soc Proc* 16: 62–65
- Gary HL (1963) Root distribution of five-stamen tamarisk, seepwillow, and arrowweed. *For Sci* 9: 311–314
- Gifford GF (1966) Aspen root studies on three sites in northern Utah. *Am Midl Nat* 75: 132–141
- Giliberto J, Estay H (1978) Seasonal water stress in some Chilean matorral shrubs. *Bot Gaz* 139: 236–260
- Ginzburg C (1966) Xerophytic structures in the roots of desert shrubs. *Ann Bot* 30: 403–418
- Gregory PJ, McGowan M, Biscoe PV, Hunter B (1978) Water relations of winter wheat. I Growth of the root system. *J Agric Sci* 91: 91–103
- Hamblin A, Tennant D (1987) Root length density and water uptake in cereals and grain legumes: how well are they correlated? *Aust J Agric Res* 38: 513–527
- Hanes TL (1965) Ecological studies on two closely related chaparral shrubs in southern California. *Ecol Monogr* 35: 213–235
- Harris GA (1967) Some competitive relationships between *Agropyron spicatum* and *Bromus tectorum*. *Ecol Monogr* 37: 89–111
- Harris WF, Kinerson RS, Edwards NT (1977) Comparison of belowground biomass of natural deciduous forest and loblolly pine plantations. *Pedobiologia* 17: 369–381
- Heitschmidt RK, Ansley RJ, Dowhower SL, Jacoby PW, Price DL (1988) Some observations from the excavation of honey mesquite root systems. *J Range Manage* 41: 227–231
- Hellmers H, Horton JS, Juhren G, O'Keefe J (1955) Root systems of some chaparral plants in southern California. *Ecology* 36: 667–678
- Heyward F (1933) The root system of longleaf pine on the deep sands of Western Florida. *Ecology* 14: 137–148
- Higgins KB, Lamb AJ, Wilgen BW van (1987) Root systems of selected plant species in mesic mountain fynbos in the Jonkershoek Valley, south-western Cape Province. *S Afr J Bot* 53: 249–257
- Hnatiuk RJ, Hopkins AJM (1980) Western Australian species-rich kwongan (sclerophyllous shrubland) affected by drought. *Aust J Bot* 28: 573–585
- Hodgkinson DC (1992) Water relations and growth of shrubs before and after fire in a semi-arid woodland. *Oecologia* 90: 467–473
- Holch AE, Hertel EW, Oakes WO, Whitwell HH (1941) Root habits of certain plants of the foothill and alpine belts of Rocky Mountain National Park. *Ecol Monogr* 11: 327–345
- Horton KW (1958) Rooting habits of lodgepole pine (Forest Research Division technical note 67). Depart North Can. Dep. North. Aff. Natl. Resourc.
- Huang JS (1987) Interactions of nematodes with rhizobia. In: Veech JA, Dickson DW (eds) *Vistas on nematology: a commemoration of the twenty-fifth anniversary of the Society of Nematologists*. Painter, DeLeon Springs, pp 301–306
- Huttel C (1975) Root distribution and biomass in three Ivory Coast rain forest plots. In: Golley FB, Medina E (eds) *Tropical ecological systems. (Ecological studies 11)*. Springer, Berlin Heidelberg New York, pp 123–130
- Ignatenko V, Khakimzyanova FI (1971) Soils and total phytomass reserves in dwarf birch – white dryas and willow tundras of the east European northlands. *Sov J Ecol* 2: 300–305
- Incoll WD (1969) Root excavation of *Eucalyptus regnans* (Research activity 69). Forest Commission, Victoria, pp 15–16
- Jaafar MN, Stone LR, Goodrum DE (1993) Rooting depth and dry matter development of sunflower. *Agron J* 85: 281–286

- Jackson RB, Canadell J, Ehleringer JR, Mooney HA, Sala OE, Schulze E-D (1996) A global analysis of root distributions for terrestrial biomes. *Oecologia* 108: 389–411
- Jeník J (1971) Root structure and underground biomass in equatorial forests. In: Duvigneaud P (ed) Productivity of forest ecosystems. Ecology and conservation 4. UNESCO Paris, pp 323–331
- Jenkins MB, Virginia RA, Jarrel WM (1988) Depth distribution and seasonal populations of mesquite-nodulating rhizobia in warm desert ecosystems. *Soil Sci Soc Am J* 52: 1644–1650
- Jennings CMH (1974) The hydrology of Botswana. PhD thesis, University of Natal, South Africa
- Joffre R, Leiva MJ, Rambal S, Fernández R. (1987) Dynamique racinaire et extraction de l'eau du sol par des graminées pérennes et annuelles méditerranéennes. *Acta Oecol Oecol Plant* 8: 181–194
- Karizumi N (1979) Illustrations of tree roots. Seibundo Shinkosha, Tokyo, pp 1121
- Kerfoot O (1963) The root system of tropical trees. *Comm For Rev* 42: 19–26
- Khodachek EA (1971) Vegetal matter of tundra phytocoenoses in the Western Part of Taimyr peninsula (International tundra biome translation 5). IBP, University of Alaska, Fairbanks
- Kimber PC (1974) The root system of jarrah (*Eucalyptus marginata*). (WA research paper 10), Forestry Department, Perth
- Klinge H (1973) Root mass estimation in lowland tropical rain forests of Central Amazonia, Brazil. II. "Coarse root mass" of trees and palms in different height classes. *An Acad Brasil Cienc* 45: 595–609
- Kochenderfer JN (1973) Root distribution under some forest types native to West Virginia. *Ecology* 54: 445–448
- Kutschera L (1960) Wurzelatlas. DLG, Frankfurt
- Lawrence T (1975) Comparison of root penetration of alтай wild ryegrass and russian wild ryegrass. *Can J Plant Sci* 55: 851–852
- Leaf AL, Leonard RE, Berglund JV (1955) Root distribution of plantation-grown red pine in an outwash soil. *Ecology* 52: 153–158
- Lewis DC, Burgy RH (1964) The relationship between oak tree roots and groundwater in fractured rock as determined by tritium tracing. *J Geophys Res* 69: 2579–2588
- Leyshon AJ (1991) Effect of rate of nitrogen fertilizer on the above- and below-ground biomass of irrigated bromegrass in southwest Saskatchewan. *Can J Plant Sci* 71: 1057–1067
- Low AB, Lamont BB (1990) Aerial and below-ground phytomass of *Banksia* scrub-heath at Aneabba, South-western Australia. *Aust J Bot* 38: 351–359
- Lull HW, Axley JH (1958) Forest soil-moisture relations in the coastal plain sands of southern New Jersey. *For Sci* 4: 2–19
- Mayaki JWC, Teare ID, Stone LR (1976) Top and root growth of irrigated and nonirrigated soybeans. *Crop Sci* 16: 92–94
- McKell CM, Jones MB, Perrier ER (1962) Root production and accumulation of root material on fertilized annual range. *Agron J* 54: 459–462
- McQuilkin WE (1935) Root development of pitch pine with some comparative observations on short-leaf pine. *J Agric Res* 51: 983–1016
- Mensah KOA, Jenik J (1968) Root system of tropical trees. 2. Features of the root system of Iroko (*Clorophora excelsa* Beth et Hook.). *Preslia Praha* 40: 21–27
- Mooney HA, Gulmon SL, Rundel PW, Ehleringer J (1980) Further observations on the water relations of *Prosopis tamarugo* of the northern Atacama desert. *Oecologia* 44: 177–180
- Nambiar EKS, Sands R (1992) Effects of compaction and simulated root channels in the subsoil on root development, water uptake and growth of radiata pine. *Tree Physiol* 10: 297–306
- Nepstad DC, de Carvalho CR, Davidson EA, Jipp PH, Lefebvre PA, Negreiros GH, da Silva ED, Stone TA, Trumbore SE, Vieira S (1994) The role of deep roots in the hydrological and carbon cycles of Amazonian forests and pastures. *Nature* 372: 666–669
- Nicoullaud B, King D, Tardieu F (1994) Vertical distribution of maize roots in relation to permanent soil characteristics. *Plant Soil* 159: 245–254
- Nilsen ET, Sharifi MR, Rundel PW, Jarrell WM, Virginia R (1983) Diurnal and seasonal water relations of the desert phreatophyte *Prosopis glandulosa* (honey mesquite) in the Sonoran desert of California. *Ecology* 64: 1381–1393
- Okali DUU, Hall JB, Lawson GW (1973) Root distribution under a thicket clump on the Accra plains, Ghana: its relevance to clump localization and water relations. *J Ecol* 61: 439–454
- Oppenheimer HR (1958) Further observations on roots penetrating into rocks and their structure. *Bull Res Counc Israel* 6: 18–31
- Orshansky G (1951) Ecological studies on lithophytes. *Palest J Bot Jerusalem* 5: 119–128
- Pages L, Pellerin S (1994) Evaluation of parameters describing the root system architecture of field grown maize plants (*Zea mays* L.). *Plant Soil* 164: 169–176
- Pate JS, Jeschke WD, Aylward MJ (1995) Hydraulic architecture and xylem structure of the dimorphic root systems of South-West Australian species of Proteacea. *J Exp Bot* 46: 907–915
- Persson H (1975) Deciduous woodland at Andersby, eastern Sweden: Field layer and below-ground production. *Acta Phytogeogr Suec* 62: 1–71
- Phillips WS (1963) Depth of roots in soil. *Ecology* 44: 424
- Plum KA, Gosting VA (1973) Origin of Australian bauxite deposits (record 1973/156). Bureau of Mineral Resources, Geology and Geophysics, Australian Department of Minerals and Energy
- Poels RLH (1987) Soils, water and nutrients in a forest ecosystem in Surinam. Agricultural University, Wageningen, The Netherlands
- Poole DK, Miller PC (1975) Water relations of selected species of chaparral and coastal sage communities. *Ecology* 56: 1118–1128
- Rawitscher F (1948) The water economy of the vegetation of the Campos cerrados in southern Brazil. *J Ecol* 36: 237–268
- Rawitscher F, Ferri MG, Rachid M (1943) Profundidade dos solos e vegetacao em campos cerrados do Brasil meridional. *Ann Acad Bras Sci* 15: 267–294
- Rees KCJ van, Comerford NB (1986) Vertical root distribution and strontium uptake of a slash pine stand on a Florida podsol. *Soil Sci Soc Am J* 50: 1042–1104
- Reicosky DC, Millington RJ, Kute A, Peters DB (1964) Patterns of water uptake and root distribution of soybeans (*Glycine max*) in the presence of a water table. *Agron J* 64: 292–297
- Reynolds TD, Fraley L Jr (1989) Root profiles of some native and exotic plant species in southeastern Idaho. *Environ Exp Bot* 29: 241–248
- Richards JH, Caldwell MM (1987) Hydraulic lift: substantial nocturnal water transport between soil layers by *Artemisia tridentata* roots. *Oecologia* 73: 486–489
- Richter DD, Makewitz D (1995) How deep is soil? *BioScience* 45: 600–609
- Roberts J (1976) A study of root distribution and growth in a *Pinus sylvestris* L. (Scots pine) plantation in East Anglia. *Plant Soil* 44: 607–621
- Rutherford MC (1983) Growth rates, biomass and distribution of selected woody plant roots in *Burkea african-Ochna pulchra* savanna. *Vegetatio* 52: 45–63
- Sala OE, Golluscio RA, Laueronh WK, Soriano A (1989) Resource partitioning between shrubs and grasses in the Patagonian steppe. *Oecologia* 81: 501–505
- Saunier RE, Wagle RF (1967) Factors affecting the distribution of shrub live oak (*Quercus turbinella* Greene). *Ecology* 48: 35–41
- Schachtschabel P, Blume HP, Hartge H, Schwertmann U (1992) *Lehrbuch der Bodenkunde*. 13th edn. Ferdinand Emke, Stuttgart
- Schulze E-D, Bauer G, Buchmann N, Canadell J, Ehleringer JR, Jackson RB, Jobbagy E, Loreti J, Mooney HA, Oesterheld M, Sala O (1996) Water availability, rooting depth, and vegetation zones along an aridity gradient in Patagonia. *Oecologia* 108: 503–511

- Schuster JL (1964) Root development of native plants under three grazing intensities. *Ecology* 45: 63–70
- Shachori A, Rosenzweig D, Poljakoff-Mayber A (1967) Effect of Mediterranean vegetation on the moisture regime. In: Sopper WE, Lull HW (eds) *Forest hydrology*. Pergamon, Oxford, pp 291–311
- Shmueli E (1948) The water balance of some plants of the Dead Sea salines. *Pal J Bot Jerusalemm Ser IV*: 117–144
- Siegenthaler U, Sarmiento JL (1993) Atmospheric carbon dioxide and the ocean. *Nature* 365: 119–125
- Silva S, Whitford WG, Jarrell WM, Virginia RA (1989) The microarthropod fauna associated with a deep rooted legume, *Prosopis glandulosa*, in the Chihuahuan desert. *Biol Fert Soils* 7: 330–335
- Specht RL, Rayson P (1957) Dark Island Heath (Ninety-Mile Plain, South Australia). III. The root systems. *Aust J Bot* 5: 103–114
- Sperry TM (1935) Root systems in Illinois prairie. *Ecology* 16: 178–202
- Stone EL, Kalisz PJ (1991) On the maximum extent of tree roots. *For Ecol Manage* 46: 59–102
- Strang RM (1969) Soil moisture relations under grassland and under woodland in the Rhodesian Highveld. *Commonw For Rev* 48: 26–40
- Strong WL, La Roi GH (1983) Root-system morphology of common boreal forest trees in Alberta, Canada. *Can J For Res* 13: 1164–1173
- Thomas CM, Davis SD (1989) Recovery patterns of three chaparral shrub species after wildfire. *Oecologia* 80: 309–320
- Tolstead WT (1942) Vegetation of the northern part of Cherry county, Nebraska. *Ecol Monogr* 12: 255–292
- Trumbore SE, Davidson EA, Barbosa PC, Nepstad DC, Martinelli LA (1995) Belowground cycling of carbon in forests and pastures of Eastern Amazonia. *Global Biogeochem Cycles* 9: 515–528
- Verzunov AI (1980) Growth of the larch and resistance of cultivated phytocenoses with its domination on semihydromorphic soils in the forest steppe in North Kazakhstan. *Sov J Ecol* 11: 98–103
- Weaver JE (1919) The Ecological relations of roots. *Carnegie Inst Wash Publ* 286: 1–128
- Weaver JE (1926) *Root development of field crops*. McGraw-Hill, New York
- Weaver JE (1958) Summary and interpretation of underground development in natural grassland communities. *Ecol Monogr* 28: 55–78
- Weaver JE, Darland RW (1949) Soil-root relationships of certain native grasses in various soil types. *Ecol Monogr* 19: 303–338
- Weaver JE, Kramer J (1932) Root system of *Quercus macrocarpa* in relation to the invasion of prairie. *Bot Gaz* 94: 51–85
- Webber PJ, May DE (1977) The magnitude and distribution of belowground plant structures in the alpine tundra of Niwot Ridge, Colorado. *Arct Alp Res* 9: 157–174
- Wein RW, Bliss LC (1974) Primary production in arctic cottongrass tussock tundra communities. *Arct Alp Res* 6: 261–274
- Welbank PJ, Gibb MJ, Taylor PJ, Williams ED (1974) Root growth of cereal crops. *Experimental Station Annual Report 1973 Part 2, Rothamsted*, pp 26–66
- Westman WE, Rogers RW (1977) Biomass and structure of a subtropical Eucalypt forest, North Stradbroke Island. *Aust J Bot* 25: 171–191
- White PD, Wood RS (1958) Growth variations in a red pine plantation influenced by a deep-lying fine soil layer. *Soil Sci Soc Proc* 22: 174–177
- Wright CD (1928) An ecological study of *Baccharis pilularis*. Ms thesis, University of California, Berkeley
- Zohary M (1961) On the hydro-ecological relations of the near east desert vegetation. In: UNESCO (ed) *Arid zone research 16*. UNESCO, Paris, pp 199–212
- Zohary M, Fahn A (1952) Ecological studies on East Mediterranean dune plants. *Bull Res Council Israel* 1: 38–53
- Zohary M, Orshan G (1949) Structure and ecology of the vegetation in the Dead Sea regions of Palestine. *Pal J Bot Jerusalem Ser IV*: 177–207
- Zwieniecki MA, Newton M (1994) Root distribution of 12-year forests at rocky sites in southwestern Oregon: effects of rock physical properties. *Can J For Res* 24: 1791–1796